

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
12 April 2001 (12.04.2001)

PCT

(10) International Publication Number
WO 01/25465 A1(51) International Patent Classification?: **C12N 15/864**,
7/01, 5/10, A61K 48/0052246 (US). ZIYING, Yan [CN/US]; Apartment 18, 733
Michael Street, Iowa City, IA 52246 (US).

(21) International Application Number: PCT/US00/27863

(74) Agent: VIKSNINS, Ann, S.; Schwegman, Lundberg,
Woessner & Kluth, P.O. Box 2938, Minneapolis, MN
55402 (US).

(22) International Filing Date: 6 October 2000 (06.10.2000)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:
60/158,209 7 October 1999 (07.10.1999) US(71) Applicant (*for all designated States except US*): UNIVER-
SITY OF IOWA RESEARCH FOUNDATION [US/US];
100 Oakdale Campus, #214 TIC, Iowa City, IA 52242-5000
(US).(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicants and


(72) Inventors: ENGELHARDT, John, F. [US/US]; 8 Laredo
Court, Iowa City, IA 52246 (US). DONGSHENG, Duan
[CN/US]; Apartment B6, 625 Emerald Street, Iowa City, IA

Published:

— With international search report.

[Continued on next page]

(54) Title: ADENO-ASSOCIATED VIRUSES AND USES THEREOF



(57) Abstract: The invention provides an isolated and purified DNA molecule comprising at least one DNA segment, a biologically active subunit or variant thereof, of a circular intermediate of adeno-associated virus, which DNA segment confers increased episomal stability, persistence or abundance of the isolated DNA molecule in a host cell. The invention also provides a composition comprising at least two adeno-associated virus vectors.

BEST AVAILABLE COPY

WO 01/25465 A1



— *Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

ADENO-ASSOCIATED VIRUSES AND USES THEREOF

Statement of Government Rights

This invention was made at least in part with a grant from the Government of the United States of America (grant HL51887 from the National
5 Institutes of Health). The Government may have certain rights in the invention.

Background of the Invention

Adeno-associated virus (AAV) is a non-pathogenic parvovirus with a single-stranded DNA genome of 4680 nucleotides. The genome may be of either
10 plus or minus polarity, and codes for two groups of genes, Rep and Cap (Berns et al., 1990). Inverted terminal repeats (ITRs), characterized by palindromic sequences producing a high degree of secondary structure, are present at both ends of the viral genome. While other members of the parvovirus group replicate autonomously, AAV requires co-infection with a helper virus (i.e.,
15 adenovirus or herpes virus) for lytic phase productive replication. In the absence of a helper virus, wild-type AAV (wtAAV) establishes a latent, non-productive infection with long-term persistence by integrating into a specific locus on chromosome 19, AAVS1, of the host genome through a Rep-facilitated mechanism (Samulski, 1993; Linden et al., 1996; Kotin et al., 1992).

20 In contrast to wtAAV, the mechanism(s) of latent phase persistence of recombinant AAV (rAAV) is less clear. rAAV integration into the host genome is not site-specific due to deletion of the AAV Rep gene (Ponnazhagan et al., 1997). Analysis of integrated proviral structures of both wild type and recombinant AAV have demonstrated head-to-tail genomes as the predominant
25 structural forms.

rAAV has recently been recognized as an extremely attractive vehicle for gene delivery (Muzyczka, 1992). rAAV vectors have been developed by substituting all viral open reading frames with a therapeutic minigene, while retaining the *cis* elements contained in two inverted terminal repeats (ITRs)
30 (Samulski et al., 1987; Samulski et al., 1989). Following transduction, rAAV genomes can persist as episomes (Flotte et al., 1994; Afione et al., 1996; Duan et al., 1998), or alternatively can integrate randomly into the cellular genome (Berns et al., 1996; McLaughlin et al., 1988; Duan et al., 1997; Fisher-Adams et al., 1996; Kearns et al., 1996; Ponnazhagan et al., 1997). However, little is

known about the mechanisms enabling rAAV vectors to persist *in vivo* or the identity of cellular factors which may modulate the efficiency of transduction and persistence. Although transduction of rAAV has been demonstrated *in vitro* in cell culture (Muzyczka, 1992) and *in vivo* in various organs (Kaplitt et al., 1994; Walsh et al., 1994; Conrad et al., 1996; Herzog et al., 1997; Snyder et al., 1997), the mechanisms of rAAV-mediated transduction remain unclear.

Moreover, while rAAV has been shown to be capable of stable, long-term transgene expression both *in vitro* and *in vivo* in a variety of tissues, the transduction efficiency of rAAV is markedly variable in different cell types. For example, rAAV has been reported to transduce lung epithelial cells at low levels (Halbert et al., 1997; Duan et al., 1998a), while high level, persistent transgene expression has been demonstrated in muscle, neurons and in other non-dividing cells (Kessler et al., 1996; Fisher et al., 1997; Herzog et al., 1997; Xiao et al., 1996; Kaplitt et al., 1994; Wu et al., 1998; Ali et al., 1996; Bennett et al., 1997 Westfall et al., 1997). These tissue-specific differences in rAAV mediated gene transfer may, in part, be due to variable levels of cellular factors affecting AAV infectivity (i.e., receptors and co-receptors such as heparin sulfate proteoglycan, FGFR-1, and $\alpha V\beta 5$ integrin) (Summerford et al., 1998; Qing et al., 1999; Summerford et al., 1999) as well as the latent life cycle (i.e., nuclear trafficking of virus and/or the conversion of single stranded genomes to expressible forms) (Qing et al., 1997; Qing et al., 1998).

Muscle-mediated gene transfer represents a very promising approach for the treatment of hereditary myopathies and several other metabolic disorders. Previous studies have demonstrated remarkably efficient and persistent transgene expression to skeletal muscle *in vivo* with rAAV vectors. Applications in this model system include the treatment of several inherited disorders such as Factor IX deficiency in hemophilia B and epo deficiencies (Kessler et al., 1996; Herzog et al., 1997). Although the conversion of low-molecular-weight rAAV genomes to high-molecular-weight concatamers has been inferred as evidence for integration of proviral DNA in the host genome, no direct evidence exists in this regard (Xiao et al., 1996; Clark et al., 1997; Fisher et al. 1997). Also, the molecular processes and/or structures associated with episomal long-term persistence of rAAV genomes, e.g., in nondividing mature myofibers, remains

unclear.

In addition, due to limitations in rAAV vector packaging capacity, a rAAV vector may not be useful if large regulatory elements are needed to control transgene expression.

- 5 Thus, there is a need for rAAV vectors that have increased stability and/or persistence in host cells. Moreover, there is a need for vectors useful to express large open reading frames.

Summary of the Invention

- 10 The present invention provides a recombinant adeno-associated virus (rAAV) vector comprising a nucleic acid segment formed by the juxtaposition of sequences in the AAV inverted terminal repeats (ITRs) which are present in a circular intermediate of AAV. The circular intermediate was isolated from rAAV-infected cells by employing a recombinant AAV "shuttle" vector. The
- 15 shuttle vector comprises: a) a bacterial origin of replication; b) a marker gene or a selectable gene; c) a 5' ITR; and d) a 3' ITR. Preferably, the recombinant AAV shuttle vector contains a reporter gene, e.g., a GFP, alkaline phosphatase or β -galactosidase gene, a selectable marker gene, e.g., an ampicillin-resistance gene, a bacterial origin of replication, a 5' ITR and a 3' ITR. The vector is
- 20 contacted with eukaryotic cells so as to yield transformed eukaryotic cells. Low molecular weight DNA ("Hirt DNA") from the transformed eukaryotic cells is isolated. Bacterial cells are contacted with the Hirt DNA so as to yield transformed bacterial cells. Then bacterial cells are identified which express the marker or selectable gene present in the shuttle vector and which comprise at
- 25 least a portion of a circular intermediate of adeno-associated virus. Also, as described below, it was found that circularized intermediates of rAAV impart episomal persistence to linked sequences in HeLa cells, fibroblasts and muscle cells. In HeLa cells, the incorporation of certain AAV sequences, e.g., ITRs, from circular intermediates into a heterologous plasmid conferred a 10-fold
- 30 increase in the stability of plasmid-based vectors in HeLa cells. Unique features of these transduction intermediates included the *in vivo* circularization of a head-to-tail monomer as well as multimer (concatamers) episomal viral genomes with associated specific base pair alterations in the 5' viral D-sequence. The

majority of circular intermediates had a consistent head-to-tail configuration consisting of monomer genomes (<3 kb) which slowly converted to large multimers of >12 kb by 80 days post-infection in muscle. Importantly, long-term transgene expression was associated with prolonged (80 day) episomal persistence of these circular intermediates. Thus, *in vivo* persistence of rAAV can occur through episomal circularized genomes which may represent prointegration intermediates with increased episomal stability. Moreover, as described below, co-infection with adenovirus, at high multiplicities of infection (MOI) capable of producing early adenoviral gene products, led to increases in the abundance and stability of AAV circular intermediates which correlated with an elevation in transgene expression from rAAV vectors. Thus, these results demonstrate the existence of a molecular structure involved in AAV transduction which may play a role in episomal persistence and/or integration.

Further, these results may aid in the development of non-viral or viral-based gene delivery systems having increased efficiency. For example, therapeutic or prophylactic therapies in which the present vectors are useful include blood disorders (e.g., sickle cell anemia, thalassemias, hemophilias, and Fanconi anemias), neurological disorders, such as Alzheimer's disease and Parkinson's disease, and muscle disorders involving skeletal, cardiac or smooth muscle. In particular, therapeutic genes useful in the vectors of the invention include the β -globin gene, the γ -globin gene, the cystic fibrosis transmembrane conductance receptor gene (CFTR), the erythropoietin (epo) gene, the Fanconi anemia complementation group, a gene encoding a ribozyme, an antisense gene, a low density lipoprotein (LDL) gene, a tyrosine hydroxylase gene (Parkinson's disease), a glucocerebrosidase gene (Gaucher's disease), an arylsulfatase A gene (metachromatic leukodystrophies) or genes encoding other polypeptides or proteins. Also within the scope of the invention is the inclusion of more than one gene in a vector of the invention, i.e., a plurality of genes may be present in an individual vector. Further, as a circular intermediate may be a concatamer, each monomer of that concatamer may comprise a different gene, or a portion thereof.

For viral-based delivery systems, helper-free virus can be prepared (see WO 95/13365) from circular intermediates or vectors of the invention.

Alternatively, liposomes, plasmid or virosomes may be employed to deliver a vector of the invention to a host or host cell.

The increased persistence of circular intermediates or vectors having one or a plurality of ITRs may be due to the primary and/or secondary structure of the ITRs. The primary structure of a consensus sequence (SEQ ID NO:3) of ITRs formed by the juxtaposition and physical (phosphodiester bond) linkage of ITRs from AAV is shown in Figure 2C. However, as described hereinbelow, each ITR sequence may be incomplete, i.e., the ITR may be a subunit or portion of the full length ITRs present in the consensus sequence. Moreover, preferably, an isolated DNA segment of the invention is not the 165 bp double DD sequence (SEQ ID NO:7) disclosed in U.S. Patent No. 5,478,745, referred to as a "double sequence".

Moreover, the formation, persistence and/or abundance of molecules having the ITR sequences of the invention may be modulated by helper virus, e.g., adenoviral proteins and/or host cell proteins. Thus, the circular intermediates or vectors of the invention may be useful to identify and/or isolate proteins that bind to the ITR sequences present in those molecules.

Therefore, the present invention provides an isolated and purified DNA molecule comprising at least one DNA segment, a biologically active subunit or variant thereof, of a circular intermediate of adeno-associated virus, which DNA segment confers increased episomal stability, persistence or abundance of the isolated DNA molecule in a host cell. Preferably, the DNA molecule comprises at least a portion of a left (5') inverted terminal repeat (ITR) of adeno-associated virus. Also preferably, the DNA molecule comprises at least a portion of a right (3')-inverted terminal repeat of adeno-associated virus. The invention also provides a gene transfer vector, comprising: at least one first DNA segment, a biologically active subunit or variant thereof, of a circular intermediate of adeno-associated virus, which DNA segment confers increased episomal stability or persistence of the vector in a host cell; and a second DNA segment comprising a gene. Preferably, the second DNA segment encodes a therapeutically effective polypeptide. The first DNA segment comprises ITR sequences, preferably at least about 100, more preferably at least about 300, and even more preferably at least about 400, bp of adeno-associated virus sequence. A preferred vector of the

invention is a plasmid.

Thus, the vector of the invention is useful in a method of delivering and/or expressing a gene in a host cell, to prepare host cells having the vector(s), and in the preparation of compositions comprising such vectors. To deliver the
5 gene to the host cell, a recombinant adenovirus helper virus may be employed.

The implications of intermolecular recombination of rAAV genomes to form a single circular episome, which may be a circular concatamer comprising at least two different rAAV genomes, is particularly relevant for gene therapy with rAAV. First, large regulatory elements and genes beyond the packaging
10 capacity of rAAV can be brought together by co-infecting tissue with two independent vectors. For example, enhancers and/or promoters may be introduced into one vector while DNA comprising an open reading frame, e.g., a gene of interest, with or without a minimal promoter, is introduced into a second vector. Thus, after co-infection with the two vectors, the transgene cassette size
15 is increased beyond that for a single AAV vector alone and the DNA comprising the opening reading frame is linked to the enhancer and/or promoter. In another embodiment, of the invention, vectors encoding two independent regions of a gene are brought together to form an intact splicing unit by circular concatamerization. In a further embodiment of the invention, a vector
20 comprising an origin of replication and a DNA encoding a protein that binds to the origin and promotes replication and/or maintenance of DNA that is linked to the origin, and a vector comprising a gene of interest are brought together after co-infection to form an autonomously replicating episome comprising the gene.

As described hereinbelow, the tibialis muscle of mice was co-infected
25 with rAAV Alkaline phosphatase (Alkphos) and GFP encoding vectors. The GFP shuttle vector also encoded ampicillin resistance and a bacterial origin of replication to allow for bacterial rescue of circular intermediates in Hirt DNA from infected muscle samples. There was a time dependent increase in the abundance of rescued plasmids encoding both GFP and Alkphos that reached
30 33% of the total circular intermediates by 120 days post-infection. Furthermore, these large circular concatamers were capable of expressing both GFP and Alkphos encoded transgenes following transient transfection in cell lines. Thus, concatamerization of AAV genomes *in vivo* occurs through intermolecular

recombination of independent monomer circular viral genomes. Therefore, a plurality of DNA segments, each in an individual rAAV vector, may be delivered so as to result in a single DNA molecule having a plurality of the DNA segments. For example, one rAAV vector comprises a first DNA segment
5 comprising a 5' ITR linked to a second DNA segment comprising a promoter operably linked to a third DNA segment comprising a first open reading frame linked to a fourth DNA segment comprising a 3' ITR. A second rAAV vector comprises a first DNA segment comprising a 5' ITR linked to a second DNA segment comprising a promoter operably linked to a third DNA segment
10 comprising a second open reading frame linked to a fourth DNA segment comprising a 3' ITR.

In another embodiment, one rAAV vector comprises a first DNA segment comprising a 5' ITR linked to a second DNA segment comprising a promoter operably linked to a third DNA segment comprising the 5' end of an
15 open reading frame linked to fourth DNA segment comprising a 5' splice site linked to a fifth DNA segment comprising a 3' ITR. The second rAAV vector comprises a first DNA segment comprising a 5' ITR linked to a second DNA segment comprising a 3' splice site linked to a third DNA segment comprising the 3' end of the open reading frame linked to a fourth DNA segment comprising
20 a 3' ITR. Preferably, the second and third DNA segments together comprise DNA encoding, for example, CFTR, factor VIII, dystrophin, or erythropoietin. Also preferably, the second DNA segment comprises the endogenous promoter of the respective gene, e.g., the epo promoter.

Thus, the invention provides a composition comprising: a first adeno-
25 associated virus vector comprising linked DNA segments and at least a second adeno-associated virus comprising linked DNA segments. The linked DNA segments of the first vector comprise: a first DNA segment comprising a 5' ITR; a second DNA segment comprising at least a portion of an open reading frame operably linked to a promoter, wherein the DNA segment does not
30 comprise the entire open reading frame; a third DNA segment comprising a splice donor site; and iv) a fourth DNA segment comprising a 3' ITR. The linked DNA segments of the second vector comprise a first DNA segment comprising a 5' ITR; a second DNA segment comprising a splice acceptor site; a

third DNA segment comprising at least a portion of an open reading frame which together with the second DNA segment of the first vector encodes a full-length polypeptide; and a fourth DNA segment comprising a 3' ITR. Preferably, the second DNA segment of the first vector comprises a first exon of a gene

5 comprising more than one exon and the third DNA segment of the second vector comprises at least one exon of a gene that is not the first exon.

The invention also provides a method to transfer and express a polypeptide in a host cell. The method comprises contacting the host cell with at least two rAAV vectors. One rAAV vector comprises a first DNA segment

10 comprising a 5' ITR linked to a second DNA segment comprising a promoter operably linked to a third DNA segment comprising a first open reading frame linked to a fourth DNA segment comprising a 3' ITR. A second rAAV vector comprises a first DNA segment comprising a 5' ITR linked to a second DNA segment comprising a promoter operably linked to a third DNA segment

15 comprising a second open reading frame linked to a fourth DNA segment comprising a 3' ITR. Alternatively, one rAAV vector comprises a first DNA segment comprising a 5' ITR linked to a second DNA segment comprising a promoter operably linked to a third DNA segment comprising the 5' end of an open reading frame linked to fourth DNA segment comprising a 5' splice site

20 linked to a fifth DNA segment comprising a 3' ITR. The second rAAV vector comprises a first DNA segment comprising a 5' ITR linked to a second DNA segment comprising a 3' splice site linked to a third DNA segment comprising the 3' end of the open reading frame linked to a fourth DNA segment comprising a 3' ITR. The host cell is preferably contacted with both of the vectors,

25 concurrently, although it is envisioned that the host cell may be contacted with each vector at a different time relative to the contact with the other vector(s).

Also provided is a method in which the composition of the invention is administered to the cells or tissues of an animal. For example, rAAV vectors have shown promise in transferring the CFTR gene into airway epithelial cells of

30 animal models and nasal sinus of CF patients. However, high level expression of CFTR has not been achieved due to the fact that AAV cannot accommodate the full-length CFTR gene together with a potent promoter. A number of studies have tried to optimize rAAV-mediated CFTR expression by utilizing truncated

or partially deleted CFTR genes together with stronger promoters. However, it is currently unknown what effect deletions within the CFTR gene may have on complementation of bacterial colonization defects in the CF airway. Therefore, the present invention includes the administration to an animal of a composition
5 of the invention comprising at least two rAAV vectors which together encode CFTR. The present invention is useful to overcome the current size limitation for transgenes within rAAV vectors, and allows for the incorporation of a larger transcriptional regulatory region, e.g., a stronger heterologous promoter or the endogenous CFTR promoter.

10 As described hereinbelow, transgene expression from rAAV luciferase vectors, with or without a promoter, can be greatly enhanced by co-infection with an independent rAAV vector carrying the cytomegalovirus (CMV) and simian virus 40 (SV40) enhancers. Thus, co-infection with a transgene containing vector and a second vector comprising at least one, preferably at least
15 two or more, enhancer sequences, of cell lines and muscle *in vivo* resulted in a greater than 600-fold enhancement of transgene expression from a minimal SV40 promoter. Furthermore, 200-fold enhancement was also achieved by *cis*-activation of ITRs in transgene containing vectors without a promoter. Thus, large regulatory elements including tissue specific enhancers can be introduced
20 into cells by a separate rAAV to regulate the expression of a second transgene containing vector in *cis* following intracellular concatamerization.

Thus, the invention provides a composition comprising at least two recombinant AAV genomes. The composition comprises a first recombinant AAV comprising a first recombinant DNA molecule comprising linked: i) a first
25 DNA segment comprising a 5'-inverted terminal repeat of AAV; ii) a second DNA segment which does not comprise AAV sequences; and iii) a third DNA segment comprising a 3'-inverted terminal repeat of AAV; and comprises a second recombinant AAV comprising a second recombinant DNA molecule comprising linked: i) a first DNA segment comprising a 5'-inverted terminal
30 repeat of AAV; ii) a second DNA segment which does not comprise AAV sequences and which second DNA segment is different than the second DNA segment of the first recombinant DNA molecule; and iii) a third DNA segment comprising a 3'-inverted terminal repeat of AAV. The composition of the

invention, is preferably contacted with a mammalian host cell, e.g., a murine, canine, feral or human cell. Alternatively, a host cell may be contacted with each recombinant AAV individually, e.g., sequentially.

Thus, in one embodiment of the invention, a host cell is contacted with at least two recombinant AAV genomes. A first recombinant AAV comprises a first recombinant DNA molecule comprising linked: i) a first DNA segment comprising a 5'-inverted terminal repeat of AAV; ii) a second DNA segment which does not comprise AAV sequences; and iii) a third DNA segment comprising a 3'-inverted terminal repeat of AAV. A second recombinant AAV comprises a second recombinant DNA molecule comprising linked: i) a first DNA segment comprising a 5'-inverted terminal repeat of AAV; ii) a second DNA segment which does not comprise AAV sequences and which second DNA segment is different than the second DNA segment of the first recombinant DNA molecule; and iii) a third DNA segment comprising a 3'-inverted terminal repeat of AAV.

In one embodiment of the invention, the second DNA segment of the first recombinant DNA molecule comprises a portion of an open reading frame, e.g., an exon of a multi-exon gene, operably linked to a promoter. For example, the promoter may be the endogenous promoter for the gene corresponding to the open reading frame. Preferably, the second DNA segment of the second recombinant DNA molecule comprises the remainder of the open reading frame which together with the second DNA segment of the first recombinant DNA molecule encodes a full-length polypeptide. Also preferably, the first recombinant DNA molecule comprises a splice donor site 3' to the open reading frame, and the second DNA segment of the second recombinant DNA molecule comprises a splice acceptor site 5' to the remainder of the open reading frame.

In another embodiment of the invention, the second DNA segment of the first recombinant DNA molecule comprises at least one heterologous enhancer and/or at least one heterologous promoter, i.e., the enhancer and/or promoter sequences are not derived from AAV sequences. Preferably, the second DNA segment of the second recombinant DNA molecule comprises at least a portion of an open reading frame.

In yet a further embodiment of the invention, the second DNA segment of the first recombinant DNA molecule comprises an origin of replication functional in a host cell, e.g., a viral origin of replication such as OriP.

Preferably, the origin is functional in a human cell. Also preferably, the second DNA segment of the first recombinant DNA molecule further comprises DNA encoding a protein that binds to the origin of replication, e.g., EBNA-1. The second DNA segment in the second recombinant DNA molecule comprises at least a portion of an open reading frame, and preferably a promoter operably linked to the open reading frame.

10

In yet another embodiment of the invention, the second DNA segment of the first recombinant DNA molecule comprises a *cis*-acting integration sequence(s) for a recombinase and also encodes a recombinase or integrase that is specific for the integration sequence(s), e.g., Cre/lox system of bacteriophage P1 (U.S. Patent No. 5,658,772), the FLP/FRT system of yeast, the Gin recombinase of phage Mu, the Pin recombinase of *E. coli*, the R/RS system of the pSR1 plasmid, a retrotransposase or the integrase from a lentivirus or retrovirus. The second DNA segment in the second recombinant DNA molecule comprises at least a portion of an open reading frame, and preferably a promoter operably linked to the open reading frame. The formation of a concatamer comprising the first and the second recombinant DNA molecules, and the expression of the recombinase or integrase, will enhance the integration of the concatamer, or a portion thereof, into the host genome.

20

25

Brief Description of the Figures

Figure 1. Structure of proviral shuttle vector and the predicted structure of rAAV circular intermediate monomers. With the aid of a rAAV *cis*-acting plasmid, pCisAV.GFP3ori (Panel A), AV.GFP3ori recombinant virus was produced (Panel B). This vector encoded a GFP transgene cassette, an ampicillin resistance gene (*amp*), and a bacterial replication origin (*ori*). The predominant form of circular intermediates isolated following transduction of Hela cells with AV.GFP3ori consisted of head-to-tail monomers (Panels C and D).

30

Figure 2. Structural analysis of rAAV circular intermediates in Hela cells. Circular rAAV intermediate clones isolated from AV.GFP3ori infected Hela cells were analyzed by diagnostic restriction digestion with AseI, SphI, and PstI together with Southern blotting against ITR, GFP, and Stuffer ³²P-labeled probes. In panel A, four clones representing the diversity of intermediates found (p190, p333, p280, and p345) gave a diagnostic PstI (P) restriction pattern (3 kb and 1.7 kb bands) consistent with a circular monomer or multimer intact genome [agarose gel (Left) and Southern blot (Right)]. SphI (S) digestion demonstrated existence of a single ITR (p190), two ITRs in a head-to-tail orientation (p333 and p280), and three ITRs (p345) in isolated circular intermediates. The restriction pattern of pCisAV.GFP3ori (U; uncut, P; PstI cut, S; SphI cut) and 1 kb DNA ladder (L) are also given for comparison. One additional circular form (p340) was repetitively seen and had an unidentifiable structure which lacked intact ITR sequences. Circular concatamers were identified by partial digestion with AseI for clones p280 (dimer) and p333 (monomer) as is shown in Panel B. Sequence analysis (Panel C) of six clones (SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, and SEQ ID NO:13) with identical restriction patterns to p333 (Panel A) was performed using primers (indicated by arrows) juxtaposed to the partial p5 promoter (dotted line) and ITRs (solid line). The top sequence (SEQ ID NO:3) represents the proposed head-to-tail structure of intact ITR arrays with alignment of sequence derived from individual clones. The junction of the inverted ITRs is marked by inverted arrowheads (at 251 bp). Several consistent bp changes (shaded) were noted in the 5'ITR D-sequence (boxed) within four clones (p79, p81, p87, and p88). All bp changes are indicated in lower case letters.

Figure 3. Adenovirus augments AAV circular intermediate formation in Hela cells. Infection of Hela cells with increasing doses (0, 500, and 5000 particles/cell) of recombinant E1-deleted adenovirus (Ad.CMVlacZ) leads to substantial expression of E2a 72kd DNA Binding Protein, as demonstrated by indirect immunofluorescent staining for DBP at 72 hours post-infection (Panel A). Co-infection of Hela cells with Ad.CMVlacZ (5000 particles/cell) and AV.GFP3ori (1000 DNA particles/cell) led to substantial augmentation of rAAV GFP transgene expression (Panel B). Augmentation in rAAV GFP transgene

expression in the presence of increasing amounts (0, 500, 5000 and 10000 particles/cell) of recombinant Ad.CMVlacZ was quantified by FACS analysis at 72 hour post-infection (Panel C). Results demonstrate the mean (+/-SEM) for two experiments performed in duplicate. In addition, an aliquot of cells was split
5 (1:10) at the time of FACS analysis and GFP colony forming units (CFU) per 10X field were quantified at 6 days (CPE denotes significant cytopathic effects at an adenoviral MOI of 10,000 particles/cell and was not quantified for GFP colonies). Hirt DNAs from AV.GFP3ori (1000 DNA particles/cell) infected Hela cells with or without co-infection with Ad.CMVlacZ (5,000 particles/cell)
10 were used to transform *E. coli*. The total number of ampicillin-resistant bacterial CFU (Panel D) and total number of head-to-tail circular intermediates CFU (Panel E) are given for a representative experiment. Greater than 20 clones for each time point were evaluated by Southern blot (see Figure 2 for detail). Zero hour controls were performed by mixing an equivalent amount of AV.GFP3ori
15 virus as used in experiments with mock infected cellular lysates prior to Hirt purification. Panel F depicts the abundance of head-to-tail circular intermediates as a percentage of total ampicillin-resistant bacterial CFU isolated from Hirt DNA.

Figure 4. Formation of rAAV head-to-tail circular intermediates
20 following *in vivo* transduction of muscle. The tibialis anterior muscle of 4-5 week old C57BL/6 mice were infected with AV.GFP3ori (3 X 10¹⁰ particles) in HEPES buffered saline (30 μ l). GFP expression (Panel A) was analyzed by direct immunofluorescence of freshly excised tissues and/or in formalin-fixed cryopreserved tissue sections in four independently injected muscles harvested at
25 0, 5, 10, 16, 22 and 80 days post-infection. GFP expression was detected at low levels beginning at 10 days and was maximum at 22 days post-infection. Expression remained stable to 80 days at which time greater than 50% of the tissue was positive (see 80 day tissue cross section counter stained with propidium iodide, panel A). Hirt DNA was isolated from muscle samples at
30 each of the various time points and after points was used to transform *E. coli*. Rescued plasmids (p439, p16, p17) were analyzed by Southern blotting in Panel B showing an agarose gel on left and ITR probed blot on right. U:uncut, P:PstI cut, and S:SphI cut. The schematic drawing of the most predominant type of

head-to-tail circular AAV intermediate plasmids rescued from bacteria is given in the right of Panel B and shows the structure of p17 as an example. Other typical clones included those with less than two ITRs as shown for p16. SphI digestion of p16 and p17 plasmids released ITR hybridizing fragments of approximately 140 and 300 bp, respectively. The slightly lower mobility then predicted for these ITR fragments likely represents anomalous migration due to the high secondary structure of inverted repeats within ITRs. Sequence analysis of p17 and p16 using nested primers to 5' and 3'-ITRs also confirmed the ITR orientations shown to the right of the gel. Additional restriction enzyme analyses to determine this structure included double and single digests with SphI, PstI, AseI, and/or SmaI. An example of an atypical clone (p439) rescued from bacteria with unknown structure is also shown.

Figure 5. Frequency of circular intermediate formation in muscle following transduction with rAAV. Hirt DNAs isolated from rAAV infected tibialis muscle were used to transform *E. coli* and the rescued plasmids analyzed by Southern blotting (greater than 20 clones were analyzed from at least two independent muscle samples for each time point). The averages of total head-to-tail circular intermediate clones (line) and ampicillin resistant bacterial clones (bar) isolated from each tibialis anterior muscle at 0, 5, 10, 16, 22 and 80 days post-infection are summarized in Panel A. Only plasmids which contained 1-2 ITRs were included in the estimation of total head-to-tail circular intermediates. Plasmids which demonstrated an absence of ITR hybridizing SphI fragments (between 150 to 300 bp) were omitted from the calculations. Panel B demonstrates the diversity of ITR arrays found in head-to-tail circular intermediates at 80 days post-infection. This panel depicts a Southern blot probed with ITR sequences and represents circular intermediates with 1-3 ITRs. SphI fragments which hybridize to ITR probes indicate the size of inverted ITR arrays (marked by arrows to right of gel). Additional restriction enzyme analysis was used to determine the structure of monomer and multimer circular intermediates. Examples are shown for two multimer (p136 and p143) circular intermediates which contain approximately three AAV genomes. Undigested plasmids of p136 and p143 migrate greater than 12 kb and is contrasted to the most predominant form of head-to-tail undigested circular intermediates at 22

days which migrate at 2.5 kb. The digestion pattern of p136 is consistent with a uniform head-to-tail configuration of three genomes which is indistinguishable from digestion patterns of p139 which contains one circularized genome (undigested p139 migrates at 2.5 kb, data not shown, also see examples p17 in Figure 4). In contrast, p136 depicts a more complex head-to-tail multimer circular intermediate which has various deletions and duplications within the ITR arrays. Predicted structure of five representative intermediates is schematically shown in Panel C.

Figure 6. Molecular size of circular intermediates in muscle. Hirt DNA from AV.GFP3ori infected muscle was size fractionated by electrophoresis and various molecular weight fractions transformed into *E. coli*. Results demonstrate the abundance of circular intermediates at each of the given molecular weights at 22 and 80 days post-infection with the rAAV shuttle vector. Structure of circular intermediates were confirmed by Southern blot restriction analysis.

Figure 7. Head-to-tail circular intermediates demonstrate increased stability of GFP expression following transient transfection in Hela cells. Subconfluent monolayers of Hela cells were co-transfected with p81, p87, or pCMVGFP and pRSVlacZ as an internal control for transfection efficiency as described in the methods. Panel A demonstrates the expansion of GFP clones after one passage (arrows). Quantification of clone size and numbers are shown in Panel B. Clone size represents the mean raw values while clone numbers are normalized for transfection efficiency as determined by X-gal staining for pRSVlacZ. The data at the top of bar graph values for each construct in Panel B represents quantification of GFP clones after second passage (also normalized for transfection efficiency). Results indicate the mean (+/-SEM) of duplicate experiments with greater than 20 fields quantified for each experimental point. The persistence of transfected p81 and pCMVGFP plasmid DNA at passage-7 post-transfection was evaluated by genomic Southern blot of total cellular DNA hybridized against ³²P-labeled GFP probe (Panel C, results from two independent transfections are shown). U:uncut, C:PstI cut. The migration of uncut dimer and monomer plasmids forms are marked on the left. PstI digestion of the plasmids results in bands at 4.7 kb (pCMVGFP, single PstI site in plasmid) and 1.7 kb (p81, two PstI sites flanking the GFP gene). To determine whether the

head-to-tail ITR array within circular intermediates was responsible for increases in the persistence of GFP expression, the head-to-tail ITR DNA element was subcloned into the pGL3 luciferase plasmid to generate pGL3(ITR). Results in Panel D compare the extent of luciferase transgene expression following

5 transfection with pGL3 and pGL3(ITR) at 10 days (passage-2) post-transfection. Results are the mean (+/-SEM) for triplicate experiments and are normalized for transfection efficiency using a dual renilla luciferase reporter vector (pRLSV40, Promega).

Figure 8. Identification of adenoviral genes responsible for augmentation

10 of AAV circular intermediate formation. HeLa cells were infected with AV.GFP3ori (1000 DNA particles/cell) in the presence of wtAd5, *dl802* (E2a-deleted), and *dl1004* (E4-deleted) adenovirus (at the indicated MOIs). Total number of head-to-tail circular intermediates from Hirt DNA and the level of augmentation of GFP transgene expression (as determined by FACS) was

15 quantified at 24 hours post-infection (Panel A). Results are the average of duplicate experiments. Panel B depicts results from Southern blot analysis of Hirt DNA following hybridization to a GFP P³²-labeled probe. DNA loads were 10% of the total Hirt yield from a 35 mm plate of HeLa cells. Infections were carried out identically to that described for Panel A. Arrows mark replication

20 form concatamers (Rf_c), dimers (Rf_d), monomers (Rf_m), and single-stranded AAV genomes (ssDNA).

Figure 9. Model for independent mechanistic interactions of adenovirus with lytic and latent phase aspects of the AAV life cycle. The adenoviral E4 gene has been shown to augment the level of rAAV second strand synthesis

25 giving rise to replication form dimers (Rf_d) and monomers (Rf_m) (Figure 8B). This augmentation leads to substantial increases in transgene expression from rAAV vectors and most closely mirrors lytic phase replication of wtAAV as head-to-head and tail-to-tail concatamers. In contrast, E4 expression inhibits the formation of head-to-tail circular intermediates of AAV. Hence, it appears that

30 increases in the amount of Rf_d and Rf_m double stranded DNA genomes does not increase the extent of circular intermediate formation. Such findings suggest that conversion of Rf_m and Rf_d to circular intermediates does not likely occur and implicates two mechanistically distinct pathway for their formation. In support

of this hypothesis, adenoviral E2a gene expression does not enhance the formation of Rf_m and Rf_d genomes but rather increase the abundance and/or stability of head-to-tail circular intermediates. Furthermore, in the absence of E4, E2a gene expression does not lead to augmentation of rAAV transgene
5 expression. Since circular intermediates have increased episomal stability in muscle and in HeLa cells, this molecular structure may be important in the latent phase of AAV persistence. Alternatively, these circular intermediates may represent pre-integration complexes as previously hypothesized for Rep facilitated integration. In the absence of Rep, circular intermediates may
10 accumulate episomally in rAAV infected cells. In summary, these findings support the notion that adenovirus may modulate both latent and lytic aspects of the AAV life cycle.

Figure 10. Individual chemical sequence of SphI fragments from p81 (A; SEQ ID NO:4), p79 (B; SEQ ID NO:5), and p1202 (C; SEQ ID NO:6) AAV
15 circular intermediates. The ends of the sequence (underlined) represent SphI restriction enzyme sites within head-to-tail circular AAV genomes cloned with the AV-GFP3ori shuttle virus.

Figure 11. Chemical sequence homology of three AAV circular intermediates with various conformations of ITR arrays (SEQ ID NO:4, SEQ ID
20 NO:5 and SEQ ID NO:6). Diversity in ITR arrays are evident from the non-conserved bases marked in lower case. The ends of the sequence (underlined) represent SphI restriction enzyme sites within head-to-tail circular AAV genomes cloned with the AV.GFP3ori shuttle virus.

Figure 12A. Palindromic repeat structure derived from chemical
25 sequencing of AAV circular intermediate isolate p81. Secondary structure of the sense strand is depicted in the top box with plasmid reference given below.

Figure 12B. Palindromic repeat structure derived from chemical sequencing of AAV circular intermediate isolate p79. Secondary structure of the sense strand is depicted in the top box with plasmid reference given below.

30 Figure 12C. Palindromic repeat structure derived from chemical sequencing of AAV circular intermediate isolate p79. Secondary structure of the sense strand is depicted in the top box with plasmid reference given below.

Figure 13. Persistence of GFP expression in developing *Xenopus* embryos microinjected with AAV circular intermediate isolate p81. The extent of GFP fluorescence in tadpoles reflects the stability of episomal or integrated microinjected plasmids. Bright field image on the left is of the p81 injected embryo. The p81 injected embryo depicts fluorescence in nearly all cells by one week post-injection. In contrast, a mosaic pattern of expression in a minority of cells in pCisAV.GFPori injected embryos. The pCisAV.GFPori plasmid contains the identical promoter sequences driving GFP gene expression and two ITRs separated by stuffer sequence. These findings demonstrate that specific structural characteristics found within AAV circular intermediates are responsible for increased persistence of transgene expression.

Figure 14. Mechanistic scheme for determining pathways for rAAV circular concatamer formation. The two independent vectors used in these studies, AV.Alkphos and AV.GFP3.ori, are shown in Panel A. Restriction sites important in the structural analysis of circular intermediates are also shown. In Panel B, a schematic representation of two potential models for circular concatamer formation is depicted, along with the methods to experimentally differentiate which of these processes is active in muscle. Following co-infection of the tibialis muscle with AV.Alkphos and AV.GFP3.ori, all subsequently rescued plasmids arise solely from circular intermediates containing AV.GFP3ori genomes. If rolling circular replication is the sole mechanism of concatamerization, only GFP expressing plasmids should be rescued. In contrast, if intermolecular recombination between independently formed monomer circular intermediates is the mechanism of concatamerization, both GFP and GFP/Alkphos expressing plasmids should be rescued.

Figure 15. Co-infection of tibialis muscle of mice with AV.Alkphos and AV.GFP3ori. Transgene expression of rAAV infected tibialis muscle was determined at 14, 35, 80 (Panels A and A'), and 120 (Panels B-D) days following co-infection with 5×10^9 DNA particles each of AV.Alkphos and AV.GFP3ori. The time course of transgene expression started around 14 days and peaked by 35-80 days. The extent of co-infection of myofibers with both Alkphos and GFP rAAV was determined in serial sections of 80 and 120 day post-infection muscle samples. Panels A-C represent GFP fluorescence of

formalin fixed, cryoprotected sections, while panels A'-C' depict the histochemical staining for Alkaline phosphatase in adjacent serial sections. A short staining time (7 minutes) was necessary to observe variation in staining levels for comparison to GFP. It was found that longer staining times

5 (30 minutes) saturated the Alkphos signal. The boxed region in panels B and B' are enlarged in panels C and C', respectively. A more precise correlation of GFP and Alkphos staining in myofibers is given in Panel D in which co-localization of GFP and Alkphos expression was examined in the same section of a 120 day post-infected sample. This was performed by photographing the GFP

10 fluorescent image prior to staining for Alkphos activity. The left panel of D shows a high power Nomarski photomicrograph of a group of myofibers (traced in red), while the corresponding GFP and Alkphos staining patterns are shown in the right panel. Photomicrographs of Alkphos staining were taken with a red filter to allow for superimposition of staining patterns with GFP fluorescence.

15 Co-expression of Alkphos and GFP is shown within myofibers as a yellow/orange color. Myofibers are marked as follows: (-) negative for both Alkphos and GFP, (*) positive for only GFP, and (+) positive for both GFP and Alkphos.

Figure 16. Rescue of circular intermediates and characterization of DNA

20 hybridization patterns. Using the ampicillin resistance gene (amp) and bacterial ori incorporated into the AV.GFP3ori vector, the extent of circular intermediate formation was assessed by rescuing amp resistant plasmids following transformation of 1/5 the isolated Hirt DNA into *E. coli* Sure cells. Twenty plasmids from each muscle sample were prepared and analyzed by slot blot

25 hybridization against GFP, Alkphos, and Amp ³²P-labeled DNA probes. A representative group demonstrating the hybridization patterns is shown in Panel A. Panel B depicts the mean (+/-SEM) number of rescued bacterial plasmids that hybridized to either GFP alone, or to both GFP and Alkphos probes, following transformation of 1/5th of the Hirt DNA. These numbers were

30 calculated from the percentage of plasmids hybridizing to GFP and/or Alkphos and the total CFU plating efficiency derived from the original transformation. In total, 3 independent muscle samples were analyzed for a total of 60 plasmids at each time point. The percentage of GFP hybridization positive rescued plasmids

that also demonstrated hybridization to Alkphos is shown in Panel C. These data demonstrate an increase in the abundance of rescued GFP/Alkphos co-encoding circular intermediates over time.

Figure 17. Transgene expression from rescued circular intermediates.

5 Rescued circular intermediate plasmids were transfected into 293 cells for assessment of their ability to express encoded transgenes. In these studies all GFP hybridization positive clones from at least two muscles were tested for each time point and scored for their ability to express GFP and Alkaline phosphatase. In total at least 40 clones were evaluated for each time point. Three patterns of
10 transgene expression were observed following transfection of these plasmids: I) no gene expression (Panel A), II) GFP expression only (Panel B), and III) GFP and Alkphos expression (Panel C). Panels A-C depict Nomarski photomicrographs (left) of GFP fluorescent fields (center) and Alkphos staining of a different field from the same culture (right). The percentage of GFP
15 hybridization positive clones that also expressed GFP is shown in Panel D. Additionally, this panel illustrates the percentage of GFP expressing clones also expressing Alkphos.

Figure 18. Structural analysis of bi-functional concatamer circular intermediates. To fully characterize the nature of GFP and Alkphos co-
20 expressing circular intermediates, detailed structural analyses were performed using restriction enzyme mapping and Southern blot hybridization with GFP, Alkphos, and ITR ³²P-labeled probes. Results from Southern blot analysis of plasmid clone #33 (Panel A) and clone #5 (Panel C) are given as representative examples of circular intermediates isolated from 80 and 35 day Hirt DNA of
25 rAAV infected muscle, respectively. Agarose gels were run in triplicate for each of these clones and Southern blot filters were hybridized with one of the three DNA probes as indicated below each autoradiogram. Molecular weights (kb) are indicated to the left of the ethidium stained agarose gel and restriction enzymes are marked on the top of each gel/filter. Panels B and D give the deduced
30 structure of plasmid clones #33 and #5, respectively, as based on Southern blot analysis. For ease of comparison with the restriction maps of the viral genomes given in Figure 14A, the position of restriction enzyme sites (kb) are marked with the indicated orientation of intact viral genomes. However, in clone #33 a

deletion occurred between the AseI and HindIII site of a head-to-tail array between AV.Alkphos and AV.GFP3ori, as reflected by a 900 bp reduction in the anticipated size of HindIII/NotI and ClaI/AseI fragments (marked by asterisks in Panel A). Furthermore, the SphI site flanking an ITR was ablated in clone #5 (bands effected by this deletion are marked by asterisks in Panel C). The deletion is not reflected in the overall concatamer since the exact region involved and/or the size of the deletion is unclear. Additionally, chemical sequence evidence of rescued circular intermediates suggests that the predominant form of ITR arrays may be in a double-D structure (i.e., one ITR flanked by two D-
10 sequence rather than two ITRs) and hence ITR arrays containing fragments may appear 147 bp shorter than indicated. However, to more easily depict the orientation of viral genomes, the position of 5' and 3' ITRs is indicated rather than representing a single ITR at these junctions.

Figure 19. Application of rAAV circular concatamers to deliver trans-
15 splicing vectors with large gene inserts. Panel A depicts two rAAV vectors encoding two halves of a cDNA (red) and flanked by splice site consensus sequences (brown). Panel B depicts one potential type of intermolecular concatamer following co-infection of cells with the independent vectors shown in panel A. Full length transgene mRNA can then be produced by splicing.
20 Panel C depicts two rAAV vectors encoding two halves of a CFTR DNA flanked by a promoter and splice donor or a splice acceptor and a poly A sequence, respectively. Panel D shows one potential type of intermolecular concatamer following co-infection of cells with the vectors in Panel C.

Figure 20. Schematic representation of the rAAV vectors used for *cis*-
25 activation.

Figure 21. Strategy for enhancing rAAV gene expression through intermolecular *cis*-activation. Two independent rAAV viruses, one encoding a transgene with or without a minimal promoter (e.g., AV.SV(P)Luc) and another harboring enhancer sequences (e.g., AV.SupEnh), were used to co-infect the
30 same tissue. Subsequent concatamerization between two rAAV vectors substantially augments expression of the transgene, due to the presence of enhancer elements within the same circularized molecule.

Figure 22. Intermolecular *cis*-activation increases rAAV mediated gene transfer in fibroblasts. Human fibroblast cells were infected with the indicated rAAV vector(s) at an moi of 1000 for each individual vector. Luciferase activity was examined at 3 days post-infection. The data represent the mean +/- SEM of 5 6 independent samples for each experimental condition.

Figure 23. Intermolecular *cis*-activation increases rAAV mediated gene transfer to muscle *in vivo*. Mouse tibialis anterior muscles were infected with the indicated rAAV vector(s) at 2×10^{10} particles per viral vector in a total volume of 30 μ l PBS. The luciferase activity in rAAV infected or mock infected (PBS) 10 muscles was examined at 30 days (Panel A) and 90 days (Panel B) post-infection. The data represent the mean +/- SEM of 6 independent muscle samples for each experimental condition. Co-administration of the AV.SupEnh vector harboring enhancer elements substantially enhanced rAAV mediated luciferase expression in muscle from both the ITR and the minimal SV40 15 promoter.

Figure 24. Viral constructs for the generation of autonomously replicating rAAV vectors as circular concatamers. Panel A depicts two rAAV constructs used to test this hypothesis. One encodes the GFP transgene (green) and the other encodes the EBNA-1 (red) and OriP (purple) sequences necessary 20 for autonomous replication. Additionally, sequences encoded within the GFP vector allow for rescue of circular intermediates in bacteria. Panel B depicts one potential type of intermolecular concatamer following co-infection of the independent vectors shown in panel A.

Figure 25. rAAV vectors used to generate a *trans*-splicing vector 25 expressing genomic epo DNA. Panel A shows a schematic of the vectors. An IRES sequence and EGFP gene are included in one of the vectors to allow for direct visualization of transgene expression. Panel B depicts a potential circular concatamer formed after co-infection. The dashed lines indicate the splicing pattern. Panel C shows the hnRNA, splicing pattern and mature mRNA 30 transcripts which result from circular concatamerization of the two vectors.

Figure 26. Production of full length Epo protein following co-infection of primary fibroblasts with two independent *trans*-splicing vectors. Confluent primary fibroblasts (8×10^5 cells) were infected with 7×10^9 particles of each

AV.Epo1 and/or AV.Epo2. Epo expression was monitored by ELISA (R & D Systems, Minneapolis, MN) by harvesting culture media 24 hours following media replacement at the indicated time points. Results are presented as the mean Epo level ($n = 3$) normalized to 1×10^5 cells.

- 5 Figure 27. Functional expression of human Epo *in vivo* using *trans*-splicing AAV vectors. Hematocrits of the C57BL/6 mice were determined at 10, 21, 35, 48, 63, 80, and 94 days following infection with 3×10^{10} particles of each independent vector either together in one tibialis muscle ($n = 6$, denoted as solid squares) or independently by administration of AV.Epo1 to the right
10 tibialis muscle and AV.Epo2 to the left tibialis muscle of the same mouse ($n = 4$, denoted by solid triangles). As a control for baseline, the serum of uninfected mice ($n = 4$, denoted by open circles) was assayed.

Detailed Description of the Invention

15 **Definitions**

As used herein, the terms "isolated and/or purified" refer to *in vitro* preparation, isolation and/or purification of a nucleic acid molecule of the invention, so that it is not associated with *in vivo* substances.

- As used herein, a DNA molecule, sequence or segment of the invention
20 preferably is biologically active. A biologically active DNA molecule of the invention has at least about 1%, more preferably at least about 10%, and more preferably at least about 50%, of the activity of a DNA molecule comprising ITR sequences from a circular intermediate of AAV, e.g., a DNA molecule comprising SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, or a
25 subunit or variant thereof. The activity of a nucleic acid molecule of the invention can be measured by methods well known to the art, some of which are described hereinbelow. For example, the presence of the DNA molecule in a recombinant nucleic acid molecule in a host cell results in episomal persistence and/or increased abundance of the recombinant molecule in those cells relative to
30 corresponding cells having a recombinant nucleic acid molecule lacking a DNA molecule of the invention.

A variant DNA molecule, sequence or segment of the invention has at least about 70%, preferably at least about 80%, and more preferably at least

about 90%, but less than 100%, contiguous nucleotide sequence homology or identity to a DNA molecule comprising ITR sequences from a circular intermediate of AAV, e.g., SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, a subunit thereof. A variant DNA molecule of the invention may include nucleotide bases not present in SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, e.g., 5', 3' or internal deletions or insertions, such as the insertion of a restriction endonuclease recognition site, so long as these bases do not substantially reduce the biological activity of the molecule. A substantial reduction in activity means a reduction in activity of greater than about 50%, preferably greater than about 90%.

I. Identification of Nucleic Acid Molecules Falling Within the Scope of the Invention

A. Nucleic Acid Molecules of the Invention

1. Sources of the Nucleic Acid Molecules of the Invention

Sources of nucleotide sequences from which the present nucleic acid molecules can be obtained include AAV infected cells, e.g., any vertebrate, preferably mammalian, cellular source.

As used herein, the terms "isolated and/or purified" refer to *in vitro* isolation of a nucleic acid, e.g., DNA molecule from its natural cellular environment, and from association with other components of the cell, such as nucleic acid or polypeptide, so that it can be sequenced, replicated, and/or expressed. For example, "isolated nucleic acid" is RNA or DNA containing greater than about 50, preferably about 300, and more preferably about 500 or more, sequential nucleotide bases that comprise a DNA segment from a circular intermediate of AAV which contains at least a portion of the 5' and 3' ITRs and the D sequence, or a variant thereof, that is complementary or hybridizes, respectively, to AAV ITR DNA and remains stably bound under stringent conditions, as defined by methods well known in the art, e.g., in Sambrook et al., 1989. Thus, the RNA or DNA is "isolated" in that it is free from at least one contaminating nucleic acid with which it is normally associated in the natural source of the RNA or DNA and is preferably substantially free of any other mammalian RNA or DNA. The phrase "free from at least one contaminating source nucleic acid with which it is normally associated" includes the case where

the nucleic acid is reintroduced into the source or natural cell but is in a different chromosomal location or is otherwise flanked by nucleic acid sequences not normally found in the source cell, e.g., in a vector or plasmid. An example of isolated nucleic acid within the scope of the invention is nucleic acid that shares
5 at least about 80%, preferably at least about 90%, and more preferably at least about 95%, sequence identity with SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6, or a subunit thereof.

As used herein, the term "recombinant nucleic acid" or "preselected nucleic acid," e.g., "recombinant DNA sequence or segment" or "preselected
10 DNA sequence or segment" refers to a nucleic acid, e.g., to DNA, that has been derived or isolated from any appropriate cellular source, that may be subsequently chemically altered *in vitro*, so that its sequence is not naturally occurring, or corresponds to naturally occurring sequences that are not positioned as they would be positioned in a genome which has not been
15 transformed with exogenous DNA. An example of preselected DNA "derived" from a source, would be a DNA sequence that is identified as a useful fragment within a given organism, and which is then chemically synthesized in essentially pure form. An example of such DNA "isolated" from a source would be a useful DNA sequence that is excised or removed from said source by chemical means,
20 e.g., by the use of restriction endonucleases, so that it can be further manipulated, e.g., amplified, for use in the invention, by the methodology of genetic engineering.

Thus, recovery or isolation of a given fragment of DNA from a restriction digest can employ separation of the digest on polyacrylamide or agarose gel by
25 electrophoresis, identification of the fragment of interest by comparison of its mobility versus that of marker DNA fragments of known molecular weight, removal of the gel section containing the desired fragment, and separation of the gel from DNA. See Lawn et al., Nucleic Acids Res., 9, 6103 (1981), and Goeddel et al., Nucleic Acids Res., 8, 4057 (1980). Therefore, "preselected
30 DNA" includes completely synthetic DNA sequences, semi-synthetic DNA sequences, DNA sequences isolated from biological sources, and DNA sequences derived from RNA, as well as mixtures thereof.

Nucleic acid molecules having base pair substitutions (i.e., variants) are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of the nucleic acid molecule.

Oligonucleotide-mediated mutagenesis is a preferred method for preparing substitution variants. This technique is well known in the art as described by Adelman et al., *DNA*, 2, 183 (1983). Briefly, AAV DNA is altered by hybridizing an oligonucleotide encoding the desired mutation to a DNA template, where the template is the single-stranded form of a plasmid or bacteriophage containing the unaltered or native DNA sequence of AAV. After hybridization, a DNA polymerase is used to synthesize an entire second complementary strand of the template that will thus incorporate the oligonucleotide primer, and will code for the selected alteration in the AAV DNA.

Generally, oligonucleotides of at least 25 nucleotides in length are used. An optimal oligonucleotide will have 12 to 15 nucleotides that are completely complementary to the template on either side of the nucleotide(s) coding for the mutation. This ensures that the oligonucleotide will hybridize properly to the single-stranded DNA template molecule. The oligonucleotides are readily synthesized using techniques known in the art such as that described by Crea et al., *Proc. Natl. Acad. Sci. U.S.A.*, 75, 5765 (1978).

The DNA template can be generated by those vectors that are either derived from bacteriophage M13 vectors (the commercially available M13mp18 and M13mp19 vectors are suitable), or those vectors that contain a single-stranded phage origin of replication as described by Viera et al., *Meth. Enzymol.*, 153, 3 (1987). Thus, the DNA that is to be mutated may be inserted into one of these vectors to generate single-stranded template. Production of the single-stranded template is described in Sections 4.21-4.41 of Sambrook et al., *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, N.Y. 1989).

Alternatively, single-stranded DNA template may be generated by denaturing double-stranded plasmid (or other) DNA using standard techniques.

For alteration of the native DNA sequence (to generate amino acid sequence variants, for example), the oligonucleotide is hybridized to the single-stranded template under suitable hybridization conditions. A DNA polymerizing enzyme, usually the Klenow fragment of DNA polymerase I, is then added to synthesize the complementary strand of the template using the oligonucleotide as a primer for synthesis. A heteroduplex molecule is thus formed such that one strand of DNA encodes the mutated form of AAV, and the other strand (the original template) encodes the native, unaltered sequence of AAV. This heteroduplex molecule is then transformed into a suitable host cell, usually a prokaryote such as *E. coli* JM101. After the cells are grown, they are plated onto agarose plates and screened using the oligonucleotide primer radiolabeled with 32-phosphate to identify the bacterial colonies that contain the mutated DNA. The mutated region is then removed and placed in an appropriate vector, generally an expression vector of the type typically employed for transformation of an appropriate host.

The method described immediately above may be modified such that a homoduplex molecule is created wherein both strands of the plasmid contain the mutations(s). The modifications are as follows: The single-stranded oligonucleotide is annealed to the single-stranded template as described above. A mixture of three deoxyribonucleotides, deoxyriboadenosine (dATP), deoxyriboguanosine (dGTP), and deoxyribothymidine (dTTP), is combined with a modified thiodeoxyribocytosine called dCTP-(α S) (which can be obtained from the Amersham Corporation). This mixture is added to the template-oligonucleotide complex. Upon addition of DNA polymerase to this mixture, a strand of DNA identical to the template except for the mutated bases is generated. In addition, this new strand of DNA will contain dCTP-(α S) instead of dCTP, which serves to protect it from restriction endonuclease digestion.

After the template strand of the double-stranded heteroduplex is nicked with an appropriate restriction enzyme, the template strand can be digested with ExoIII nuclease or another appropriate nuclease past the region that contains the site(s) to be mutagenized. The reaction is then stopped to leave a molecule that

is only partially single-stranded. A complete double-stranded DNA homoduplex is then formed using DNA polymerase in the presence of all four deoxyribonucleotide triphosphates, ATP, and DNA ligase. This homoduplex molecule can then be transformed into a suitable host cell such as *E. coli* JM101.

5 For example, a preferred embodiment of the invention is an isolated and purified DNA molecule comprising a DNA segment comprising SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, a subunit thereof or a variant thereof having nucleotide substitutions, or deletions or insertions.

II. Preparation of Molecules Useful to Practice the Methods of the Invention

10 A. Nucleic Acid Molecules

1. Chimeric Expression Cassettes

To prepare expression cassettes for transformation herein, the recombinant or preselected DNA sequence or segment may be circular or linear, double-stranded or single-stranded. Generally, the preselected DNA sequence or
15 segment is in the form of chimeric DNA, such as plasmid DNA, that can also contain coding regions flanked by control sequences which promote the expression of the preselected DNA present in the resultant cell line.

As used herein, "chimeric" means that a vector comprises DNA from at least two different species, or comprises DNA from the same species, which is
20 linked or associated in a manner which does not occur in the "native" or wild type of the species.

Aside from the preselected DNA sequences described above, a portion of the preselected DNA may serve a regulatory or a structural function. For example, the preselected DNA may itself comprise a promoter that is active in
25 mammalian cells, or may utilize a promoter already present in the genome that is the transformation target. Such promoters include the CMV promoter, as well as the SV40 late promoter and retroviral LTRs (long terminal repeat elements), although many other promoter elements well known to the art may be employed in the practice of the invention.

30 Other elements functional in the host cells, such as introns, enhancers, polyadenylation sequences and the like, may also be a part of the preselected DNA. Such elements may or may not be necessary for the function of the DNA, but may provide improved expression of the DNA by affecting transcription,

stability of the mRNA, or the like. Such elements may be included in the DNA as desired to obtain the optimal performance of the transforming DNA in the cell.

“Control sequences” is defined to mean DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotic cells, for example, include a promoter, and optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

“Operably linked” is defined to mean that the nucleic acids are placed in a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a peptide or polypeptide if it is expressed as a preprotein that participates in the secretion of the peptide or polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that the DNA sequences being linked are contiguous and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accord with conventional practice.

The preselected DNA to be introduced into the cells further will generally contain either a selectable marker gene or a reporter gene or both to facilitate identification and selection of transformed cells from the population of cells sought to be transformed. Alternatively, the selectable marker may be carried on a separate piece of DNA and used in a co-transformation procedure. Both selectable markers and reporter genes may be flanked with appropriate regulatory sequences to enable expression in the host cells. Useful selectable markers are well known in the art and include, for example, antibiotic and herbicide-resistance genes, such as *neo*, *hpt*, *dhfr*, *bar*, *aroA*, *dapA* and the like. See also, the genes listed on Table 1 of Lundquist et al. (U.S. Patent No. 5,848,956).

Reporter genes are used for identifying potentially transformed cells and for evaluating the functionality of regulatory sequences. Reporter genes which encode for easily assayable proteins are well known in the art. In general, a reporter gene is a gene which is not present in or expressed by the recipient
5 organism or tissue and which encodes a protein whose expression is manifested by some easily detectable property, e.g., enzymatic activity. Preferred genes include the chloramphenicol acetyl transferase gene (cat) from Tn9 of *E. coli*, the beta-glucuronidase gene (gus) of the *uidA* locus of *E. coli*, and the luciferase gene from firefly *Photinus pyralis*. Expression of the reporter gene is assayed at
10 a suitable time after the DNA has been introduced into the recipient cells.

The general methods for constructing recombinant DNA which can transform target cells are well known to those skilled in the art, and the same compositions and methods of construction may be utilized to produce the DNA useful herein. For example, J. Sambrook et al., Molecular Cloning: A
15 Laboratory Manual, Cold Spring Harbor Laboratory Press (2d ed., 1989), provides suitable methods of construction.

2. Transformation into Host Cells

The recombinant DNA can be readily introduced into the host cells, e.g., mammalian, bacterial, yeast or insect cells by transfection with an expression
20 vector of the invention, by any procedure useful for the introduction into a particular cell, e.g., physical or biological methods, to yield a transformed cell having the recombinant DNA stably integrated into its genome or present as an episome which can persist in the transformed cells, so that the DNA molecules, sequences, or segments, of the present invention are maintained and/or expressed
25 by the host cell.

Physical methods to introduce a preselected DNA into a host cell include calcium phosphate precipitation, lipofection, particle bombardment, microinjection, electroporation, and the like. Biological methods to introduce the DNA of interest into a host cell include the use of DNA and RNA viral
30 vectors. The main advantage of physical methods is that they are not associated with pathological or oncogenic processes of viruses. However, they are less precise, often resulting in multiple copy insertions, random integration, disruption of foreign and endogenous gene sequences, and unpredictable

expression.

As used herein, the term "cell line" or "host cell" is intended to refer to well-characterized homogenous, biologically pure populations of cells. These cells may be eukaryotic cells that are neoplastic or which have been

5 "immortalized" *in vitro* by methods known in the art, as well as primary cells, or prokaryotic cells. The cell line or host cell is preferably of mammalian origin, but cell lines or host cells of non-mammalian origin may be employed, including plant, insect, yeast, fungal or bacterial sources. Generally, the preselected DNA sequence is related to a DNA sequence which is resident in the genome of the

10 host cell but is not expressed, or not highly expressed, or, alternatively, overexpressed.

"Transfected" or "transformed" is used herein to include any host cell or cell line, the genome of which has been altered or augmented by the presence of at least one preselected DNA sequence, which DNA is also referred to in the art

15 of genetic engineering as "heterologous DNA," "recombinant DNA," "exogenous DNA," "genetically engineered," "non-native," or "foreign DNA," wherein said DNA was isolated and introduced into the genome of the host cell or cell line by the process of genetic engineering. The host cells of the present invention are typically produced by transfection with a DNA sequence in a

20 plasmid expression vector, a viral expression vector, or as an isolated linear DNA sequence.

To confirm the presence of the preselected DNA sequence in the host cell, a variety of assays may be performed. Such assays include, for example, "molecular biological" assays well known to those of skill in the art, such as

25 Southern and Northern blotting, RT-PCR and PCR; "biochemical" assays, such as detecting the presence of a polypeptide expressed from a gene present in the vector, e.g., by immunological means (immunoprecipitations, immunoaffinity columns, ELISAs and Western blots) or by any other assay useful to identify molecules falling within the scope of the invention.

30 To detect and quantitate RNA produced from introduced DNA segments, RT-PCR may be employed. In this application of PCR, it is first necessary to reverse transcribe RNA into DNA, using enzymes such as reverse transcriptase, and then through the use of conventional PCR techniques amplify the DNA. In

most instances PCR techniques, while useful, will not demonstrate integrity of the RNA product. Further information about the nature of the RNA product may be obtained by Northern blotting. This technique demonstrates the presence of an RNA species and gives information about the integrity of that RNA. The presence or absence of an RNA species can also be determined using dot or slot blot Northern hybridizations. These techniques are modifications of Northern blotting and only demonstrate the presence or absence of an RNA species.

While Southern blotting and PCR may be used to detect the DNA segment in question, they do not provide information as to whether the DNA segment is being expressed. Expression may be evaluated by specifically identifying the polypeptide products of the introduced DNA sequences or evaluating the phenotypic changes brought about by the expression of the introduced DNA segment in the host cell.

III. Dosages, Formulations and Routes of Administration

Administration of a nucleic acid molecule may be accomplished through the introduction of cells transformed with the nucleic acid molecule (see, for example, WO 93/02556), the administration of the nucleic acid molecule itself (see, for example, Felgner et al., U.S. Patent No. 5,580,859, Pardoll et al., *Immunity*, 3, 165 (1995); Stevenson et al., *Immunol. Rev.*, 145, 211 (1995); Molling, I. *Mol. Med.*, 75, 242 (1997); Donnelly et al., *Ann. N.Y. Acad. Sci.*, 772, 40 (1995); Yang et al., *Mol. Med. Today*, 2, 476 (1996); Abdallah et al., *Biol. Cell*, 85, 1 (1995)), through infection with a recombinant virus or via liposomes. Pharmaceutical formulations, dosages and routes of administration for nucleic acids are generally disclosed, for example, in Felgner et al., *supra*.

Administration of the therapeutic agents in accordance with the present invention may be continuous or intermittent, depending, for example, upon the recipient's physiological condition, whether the purpose of the administration is therapeutic or prophylactic, and other factors known to skilled practitioners. The administration of the agents of the invention may be essentially continuous over a preselected period of time or may be in a series of spaced doses. Both local and systemic administration is contemplated. When the molecules of the invention are employed for prophylactic purposes, agents of the invention are amenable to chronic use, preferably by systemic administration.

One or more suitable unit dosage forms comprising the therapeutic agents of the invention, which, as discussed below, may optionally be formulated for sustained release, can be administered by a variety of routes including oral, or parenteral, including by rectal, transdermal, subcutaneous, intravenous, intramuscular, intraperitoneal, intrathoracic, intrapulmonary and intranasal routes. The formulations may, where appropriate, be conveniently presented in discrete unit dosage forms and may be prepared by any of the methods well known to pharmacy. Such methods may include the step of bringing into association the therapeutic agent with liquid carriers, solid matrices, semi-solid carriers, finely divided solid carriers or combinations thereof, and then, if necessary, introducing or shaping the product into the desired delivery system.

When the therapeutic agents of the invention are prepared for oral administration, they are preferably combined with a pharmaceutically acceptable carrier, diluent or excipient to form a pharmaceutical formulation, or unit dosage form. The total active ingredients in such formulations comprise from 0.1 to 99.9% by weight of the formulation. By "pharmaceutically acceptable" it is meant the carrier, diluent, excipient, and/or salt must be compatible with the other ingredients of the formulation, and not deleterious to the recipient thereof. The active ingredient for oral administration may be present as a powder or as granules; as a solution, a suspension or an emulsion; or in achievable base such as a synthetic resin for ingestion of the active ingredients from a chewing gum. The active ingredient may also be presented as a bolus, electuary or paste.

Pharmaceutical formulations containing the therapeutic agents of the invention can be prepared by procedures known in the art using well known and readily available ingredients. For example, the agent can be formulated with common excipients, diluents, or carriers, and formed into tablets, capsules, suspensions, powders, and the like. Examples of excipients, diluents, and carriers that are suitable for such formulations include the following fillers and extenders such as starch, sugars, mannitol, and silicic derivatives; binding agents such as carboxymethyl cellulose, HPMC and other cellulose derivatives, alginates, gelatin, and polyvinyl-pyrrolidone; moisturizing agents such as glycerol; disintegrating agents such as calcium carbonate and sodium bicarbonate; agents for retarding dissolution such as paraffin; resorption

accelerators such as quaternary ammonium compounds; surface active agents such as cetyl alcohol, glycerol monostearate; adsorptive carriers such as kaolin and bentonite; and lubricants such as talc, calcium and magnesium stearate, and solid polyethyl glycols.

5 For example, tablets or caplets containing the agents of the invention can include buffering agents such as calcium carbonate, magnesium oxide and magnesium carbonate. Caplets and tablets can also include inactive ingredients such as cellulose, pregelatinized starch, silicon dioxide, hydroxy propyl methyl cellulose, magnesium stearate, microcrystalline cellulose, starch, talc, titanium
10 dioxide, benzoic acid, citric acid, corn starch, mineral oil, polypropylene glycol, sodium phosphate, and zinc stearate, and the like. Hard or soft gelatin capsules containing an agent of the invention can contain inactive ingredients such as gelatin, microcrystalline cellulose, sodium lauryl sulfate, starch, talc, and titanium dioxide, and the like, as well as liquid vehicles such as polyethylene
15 glycols (PEGs) and vegetable oil. Moreover, enteric coated caplets or tablets of an agent of the invention are designed to resist disintegration in the stomach and dissolve in the more neutral to alkaline environment of the duodenum.

 The therapeutic agents of the invention can also be formulated as elixirs or solutions for convenient oral administration or as solutions appropriate for
20 parenteral administration, for instance by intramuscular, subcutaneous or intravenous routes.

 The pharmaceutical formulations of the therapeutic agents of the invention can also take the form of an aqueous or anhydrous solution or dispersion, or alternatively the form of an emulsion or suspension.

25 Thus, the therapeutic agent may be formulated for parenteral administration (e.g., by injection, for example, bolus injection or continuous infusion) and may be presented in unit dose form in ampules, pre-filled syringes, small volume infusion containers or in multi-dose containers with an added preservative. The active ingredients may take such forms as suspensions,
30 solutions, or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredients may be in powder form, obtained by aseptic isolation of sterile solid or by lyophilization from solution, for constitution with a suitable

vehicle, e.g., sterile, pyrogen-free water, before use.

These formulations can contain pharmaceutically acceptable vehicles and adjuvants which are well known in the prior art. It is possible, for example, to prepare solutions using one or more organic solvent(s) that is/are acceptable
5 from the physiological standpoint, chosen, in addition to water, from solvents such as acetone, ethanol, isopropyl alcohol, glycol ethers such as the products sold under the name "Dowanol", polyglycols and polyethylene glycols, C₁-C₄ alkyl esters of short-chain acids, preferably ethyl or isopropyl lactate, fatty acid triglycerides such as the products marketed under the name "Miglyol", isopropyl
10 myristate, animal, mineral and vegetable oils and polysiloxanes.

The compositions according to the invention can also contain thickening agents such as cellulose and/or cellulose derivatives. They can also contain gums such as xanthan, guar or carbo gum or gum arabic, or alternatively polyethylene glycols, bentones and montmorillonites, and the like.

15 It is possible to add, if necessary, an adjuvant chosen from antioxidants, surfactants, other preservatives, film-forming, keratolytic or comedolytic agents, perfumes and colorings. Also, other active ingredients may be added, whether for the conditions described or some other condition.

For example, among antioxidants, t-butylhydroquinone, butylated
20 hydroxyanisole, butylated hydroxytoluene and α -tocopherol and its derivatives may be mentioned. The galenical forms chiefly conditioned for topical application take the form of creams, milks, gels, dispersion or microemulsions, lotions thickened to a greater or lesser extent, impregnated pads, ointments or sticks, or alternatively the form of aerosol formulations in spray or foam form or
25 alternatively in the form of a cake of soap.

Additionally, the agents are well suited to formulation as sustained release dosage forms and the like. The formulations can be so constituted that they release the active ingredient only or preferably in a particular part of the intestinal or respiratory tract, possibly over a period of time. The coatings,
30 envelopes, and protective matrices may be made, for example, from polymeric substances, such as polylactide-glycolates, liposomes, microemulsions, microparticles, nanoparticles, or waxes. These coatings, envelopes, and protective matrices are useful to coat indwelling devices, e.g., stents, catheters,

peritoneal dialysis tubing, and the like.

The therapeutic agents of the invention can be delivered via patches for transdermal administration. See U.S. Patent No. 5,560,922 for examples of patches suitable for transdermal delivery of a therapeutic agent. Patches for transdermal delivery can comprise a backing layer and a polymer matrix which has dispersed or dissolved therein a therapeutic agent, along with one or more skin permeation enhancers. The backing layer can be made of any suitable material which is impermeable to the therapeutic agent. The backing layer serves as a protective cover for the matrix layer and provides also a support function. The backing can be formed so that it is essentially the same size layer as the polymer matrix or it can be of larger dimension so that it can extend beyond the side of the polymer matrix or overlay the side or sides of the polymer matrix and then can extend outwardly in a manner that the surface of the extension of the backing layer can be the base for an adhesive means. Alternatively, the polymer matrix can contain, or be formulated of, an adhesive polymer, such as polyacrylate or acrylate/vinyl acetate copolymer. For long-term applications it might be desirable to use microporous and/or breathable backing laminates, so hydration or maceration of the skin can be minimized.

Examples of materials suitable for making the backing layer are films of high and low density polyethylene, polypropylene, polyurethane, polyvinylchloride, polyesters such as poly(ethylene phthalate), metal foils, metal foil laminates of such suitable polymer films, and the like. Preferably, the materials used for the backing layer are laminates of such polymer films with a metal foil such as aluminum foil. In such laminates, a polymer film of the laminate will usually be in contact with the adhesive polymer matrix.

The backing layer can be any appropriate thickness which will provide the desired protective and support functions. A suitable thickness will be from about 10 to about 200 microns.

Generally, those polymers used to form the biologically acceptable adhesive polymer layer are those capable of forming shaped bodies, thin walls or coatings through which therapeutic agents can pass at a controlled rate. Suitable polymers are biologically and pharmaceutically compatible, nonallergenic and insoluble in and compatible with body fluids or tissues with which the device is

contacted. The use of soluble polymers is to be avoided since dissolution or erosion of the matrix by skin moisture would affect the release rate of the therapeutic agents as well as the capability of the dosage unit to remain in place for convenience of removal.

5 Exemplary materials for fabricating the adhesive polymer layer include polyethylene, polypropylene, polyurethane, ethylene/propylene copolymers, ethylene/ethylacrylate copolymers, ethylene/vinyl acetate copolymers, silicone elastomers, especially the medical-grade polydimethylsiloxanes, neoprene rubber, polyisobutylene, polyacrylates, chlorinated polyethylene, polyvinyl
10 chloride, vinyl chloride-vinyl acetate copolymer, crosslinked polymethacrylate polymers (hydrogel), polyvinylidene chloride, poly(ethylene terephthalate), butyl rubber, epichlorohydrin rubbers, ethylenvinyl alcohol copolymers, ethylene-vinyloxyethanol copolymers; silicone copolymers, for example, polysiloxane-polycarbonate copolymers, polysiloxanepolyethylene oxide copolymers,
15 polysiloxane-polymethacrylate copolymers, polysiloxane-alkylene copolymers (e.g., polysiloxane-ethylene copolymers), polysiloxane-alkylenesilane copolymers (e.g., polysiloxane-ethylenesilane copolymers), and the like; cellulose polymers, for example methyl or ethyl cellulose, hydroxy propyl methyl cellulose, and cellulose esters; polycarbonates; polytetrafluoroethylene;
20 and the like.

Preferably, a biologically acceptable adhesive polymer matrix should be selected from polymers with glass transition temperatures below room temperature. The polymer may, but need not necessarily, have a degree of crystallinity at room temperature. Cross-linking monomeric units or sites can be
25 incorporated into such polymers. For example, cross-linking monomers can be incorporated into polyacrylate polymers, which provide sites for cross-linking the matrix after dispersing the therapeutic agent into the polymer. Known cross-linking monomers for polyacrylate polymers include polymethacrylic esters of polyols such as butylene diacrylate and dimethacrylate, trimethylol propane
30 trimethacrylate and the like. Other monomers which provide such sites include allyl acrylate, allyl methacrylate, diallyl maleate and the like.

Preferably, a plasticizer and/or humectant is dispersed within the adhesive polymer matrix. Water-soluble polyols are generally suitable for this

purpose. Incorporation of a humectant in the formulation allows the dosage unit to absorb moisture on the surface of skin which in turn helps to reduce skin irritation and to prevent the adhesive polymer layer of the delivery system from failing.

5 Therapeutic agents released from a transdermal delivery system must be capable of penetrating each layer of skin. In order to increase the rate of permeation of a therapeutic agent, a transdermal drug delivery system must be able in particular to increase the permeability of the outermost layer of skin, the stratum corneum, which provides the most resistance to the penetration of
10 molecules. The fabrication of patches for transdermal delivery of therapeutic agents is well known to the art.

 For administration to the upper (nasal) or lower respiratory tract by inhalation, the therapeutic agents of the invention are conveniently delivered from an insufflator, nebulizer or a pressurized pack or other convenient means of
15 delivering an aerosol spray. Pressurized packs may comprise a suitable propellant such as dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount.

20 Alternatively, for administration by inhalation or insufflation, the composition may take the form of a dry powder, for example, a powder mix of the therapeutic agent and a suitable powder base such as lactose or starch. The powder composition may be presented in unit dosage form in, for example, capsules or cartridges, or, e.g., gelatine or blister packs from which the powder
25 may be administered with the aid of an inhalator, insufflator or a metered-dose inhaler.

 For intra-nasal administration, the therapeutic agent may be administered via nose drops, a liquid spray, such as via a plastic bottle atomizer or metered-dose inhaler. Typical of atomizers are the Mistometer (Wintrop) and the
30 Medihaler (Riker).

 The local delivery of the therapeutic agents of the invention can also be by a variety of techniques which administer the agent at or near the site of disease. Examples of site-specific or targeted local delivery techniques are not

intended to be limiting but to be illustrative of the techniques available.

Examples include local delivery catheters, such as an infusion or indwelling catheter, e.g., a needle infusion catheter, shunts and stents or other implantable devices, site specific carriers, direct injection, or direct applications.

- 5 For topical administration, the therapeutic agents may be formulated as is known in the art for direct application to a target area. Conventional forms for this purpose include wound dressings, coated bandages or other polymer coverings, ointments, creams, lotions, pastes, jellies, sprays, and aerosols. Ointments and creams may, for example, be formulated with an aqueous or oily
- 10 base with the addition of suitable thickening and/or gelling agents. Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilizing agents, dispersing agents, suspending agents, thickening agents, or coloring agents. The active ingredients can also be delivered via iontophoresis, e.g., as disclosed in U.S. Patent Nos. 4,140,122;
- 15 4,383,529; or 4,051,842. The percent by weight of a therapeutic agent of the invention present in a topical formulation will depend on various factors, but generally will be from 0.01% to 95% of the total weight of the formulation, and typically 0.1-25% by weight.

- Drops, such as eye drops or nose drops, may be formulated with an
- 20 aqueous or non-aqueous base also comprising one or more dispersing agents, solubilizing agents or suspending agents. Liquid sprays are conveniently delivered from pressurized packs. Drops can be delivered via a simple eye dropper-capped bottle, or via a plastic bottle adapted to deliver liquid contents dropwise, via a specially shaped closure.

- 25 The therapeutic agent may further be formulated for topical administration in the mouth or throat. For example, the active ingredients may be formulated as a lozenge further comprising a flavored base, usually sucrose and acacia or tragacanth; pastilles comprising the composition in an inert base such as gelatin and glycerin or sucrose and acacia; and mouthwashes comprising
- 30 the composition of the present invention in a suitable liquid carrier.

 The formulations and compositions described herein may also contain other ingredients such as antimicrobial agents, or preservatives. Furthermore, the active ingredients may also be used in combination with other therapeutic

agents, for example, bronchodilators.

In particular, for delivery of a vector of the invention to a tissue such as muscle, any physical or biological method that will introduce the vector into the muscle tissue of a host animal can be employed. Vector means both a bare
5 recombinant vector and vector DNA packaged into viral coat proteins, as is well known for AAV administration. Simply dissolving an AAV vector in phosphate buffered saline has been demonstrated to be sufficient to provide a vehicle useful for muscle tissue expression, and there are no known restrictions on the carriers or other components that can be coadministered with the vector (although
10 compositions that degrade DNA should be avoided in the normal manner with vectors). Pharmaceutical compositions can be prepared as injectable formulations or as topical formulations to be delivered to the muscles by transdermal transport. Numerous formulations for both intramuscular injection and transdermal transport have been previously developed and can be used in the
15 practice of the invention. The vectors can be used with any pharmaceutically acceptable carrier for ease of administration and handling.

For purposes of intramuscular injection, solutions in an adjuvant such as sesame or peanut oil or in aqueous propylene glycol can be employed, as well as sterile aqueous solutions. Such aqueous solutions can be buffered, if desired,
20 and the liquid diluent first rendered isotonic with saline or glucose. Solutions of the AAV vector as a free acid (DNA contains acidic phosphate groups) or a pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. A dispersion of AAV viral particles can also be prepared in glycerol, liquid polyethylene glycols and mixtures
25 thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms. In this connection, the sterile aqueous media employed are all readily obtainable by standard techniques well-known to those skilled in the art.

The pharmaceutical forms suitable for injectable use include sterile
30 aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be

preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol and the like), suitable mixtures thereof, and vegetable oils.

- 5 The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of a dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal and
- 10 the like. In many cases it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by use of agents delaying absorption, for example, aluminum monostearate and gelatin.

- Sterile injectable solutions are prepared by incorporating the AAV vector
- 15 in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the sterilized active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of
- 20 sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying technique which yield a powder of the active ingredient plus any additional desired ingredient from the previously sterile-filtered solution thereof.

- For purposes of topical administration, dilute sterile, aqueous solutions
- 25 (usually in about 0.1% to 5% concentration), otherwise similar to the above parenteral solutions, are prepared in containers suitable for incorporation into a transdermal patch, and can include known carriers, such as pharmaceutical grade dimethylsulfoxide (DMSO).

- The therapeutic compounds of this invention may be administered to a
- 30 mammal alone or in combination with pharmaceutically acceptable carriers. As noted above, the relative proportions of active ingredient and carrier are determined by the solubility and chemical nature of the compound, chosen route of administration and standard pharmaceutical practice.

The dosage of the present therapeutic agents which will be most suitable for prophylaxis or treatment will vary with the form of administration, the particular compound chosen and the physiological characteristics of the particular patient under treatment. Generally, small dosages will be used initially and, if necessary, will be increased by small increments until the optimum effect under the circumstances is reached. Exemplary dosages are set out in the example below.

Since AAV has been shown to have a broad host range (for pulmonary expression) and persists in muscle, the vectors of the invention may be employed to express a gene in any animal, and particularly in mammals, birds, fish, and reptiles, especially domesticated mammals and birds such as cattle, sheep, pigs, horses, dogs, cats, chickens, and turkeys. Both human and veterinary uses are particularly preferred.

The gene being expressed can be either a DNA segment encoding a protein, with whatever control elements (e.g., promoters, operators) are desired by the user, or a non-coding DNA segment, the transcription of which produces all or part of some RNA-containing molecule (such as a transcription control element, +RNA, or anti-sense molecule).

Muscle tissue is a very attractive target for *in vivo* gene delivery and gene therapy, because it is not a vital organ and is very easy to access. If a disease is caused by a defective gene product which is required to be produced and/or secreted, such as hemophilia, diabetes and Gaucher's disease, and the like, is muscle is a good candidate to supply the gene product if the appropriate gene can be effectively delivered into the cells.

Different vectors, such as naked DNA, adenovirus and retrovirus, have been utilized to directly deliver various transgenes into muscle tissues. However, neither system can offer both high efficiency and long-term expression. For naked plasmid DNA directly delivered into muscle tissue, the efficiency is not high. There are only a few cells near the injection site that can maintain transgene expression. Furthermore, the plasmid DNA in the cells remains as non-replicating episomes, i.e., in the unintegrated form. Therefore, it will be eventually lost. For adenovirus vector, it can infect the non-dividing cells, and therefore, can be directly delivered into the mature tissues such as

muscle. However, the transgene delivered by adenovirus vectors are not useful to maintain long-term expression for the following reasons. First, since adenovirus vectors still retain most of the viral genes, they are not very safe. Moreover, the expression of those genes can cause the immune system to destroy the cells containing the vectors (see, for example, Yang et al. 1994, Proc. Natl. Acad. Sci. 91:4407-4411). Second, since adenovirus is not an integration virus, its DNA will eventually be diluted or degraded in the cells. Third, due to the immune response, adenovirus vector could not be repeatedly delivered. In the case of lifetime diseases, this will be a major limitation. For retrovirus vectors, although they can achieve stable integration into the host chromosomes, their use is very restricted because they can only infect dividing cells while a large majority of the muscle cells are non-dividing.

Adeno-associated virus vectors have certain advantages over the above-mentioned vector systems. First, like adenovirus, AAV can efficiently infect non-dividing cells. Second, all the AAV viral genes are eliminated in the vector. Since the viral-gene-expression-induced immune reaction is no longer a concern, AAV vectors are safer than Ad vectors. Thirds, AAV is an integration virus by nature, and integration into the host chromosome will stably maintain its transgene in the cells. Fourth, AAV is an extremely stable virus, which is resistant to many detergents, pH changes and heat (stable at 56°C for more than an hour). It can be lyophilized and redissolved without losing its activity. Therefore, it is a very promising delivery vehicle for gene therapy.

The invention will be further described by, but is not limited to, the following examples.

25

Example 1

Materials and Methods

Construction of rAAV Shuttle Vector.

A recombinant AAV shuttle vector (AV.GFP3ori) which contained a GFP transgene cassette, bacterial ampicillin resistance gene, and bacterial origin of replication, was generated from a *cis*-acting plasmid (pCisAV.GFP3ori). Expression of the GFP gene was directed by the CMV promoter/enhancer and SV40 poly-adenylation sequences. pCisAV.GFP3ori was constructed with

pSub201 derived ITR elements (Samulski et al., 1987) and the intactness of ITR sequences was confirmed by restriction analysis with SmaI and PvuII, and by sequencing. Recombinant AAV stocks were generated by co-transfection of pCisAV.GFP3ori and pRep/Cap together with co-infection of recombinant Ad.CMVlacZ in 293 cells (Duan et al., 1997). Following transfection of forty 150 mm plates, cells were collected at 72 hours by centrifugation and resuspended in 12 ml of buffer (10 mM Tris pH 8.0). Virus was released from cells by three cycles of freeze/thawing and passaged through a 25 gauge needle six times. Cell lysates were then treated with 1.3 mg/ml DNase I at 37°C for 30 minutes and 1% deoxycholate (g/ml final) and 0.05% trypsin (g/ml final) at 37°C for 30 minutes. Samples were then placed on ice for 10 minutes and centrifuged to remove large particulate material at 3,000 rpm for 30 minutes.

rAAV was purified by isopycnic density gradient centrifugation in CsCl ($r=1.4$) in a SW55 rotor for 72 hours at 35K. Peak fractions of AAV were combined and re-purified through two more rounds of CsCl centrifugation, followed by heating at 58°C for 60 minutes to inactivate all contaminant helper adenovirus. Typically, this preparation gave approximate AAV titers of 10^{12} DNA molecules/ml and 2.5×10^8 GFP-expressing units/ml. Recombinant viral titers were assessed by slot blot and quantified against pCisAV.GFP3ori controls for DNA particles. Functional transducing units were quantified by GFP transgene expression in 293 cells. The absence of helper adenovirus was confirmed by histochemical staining of rAAV infected 293 cells for beta-galactosidase, and no recombinant adenovirus was found in 10^{10} particles of purified rAAV stocks. The absence of significant wtAAV contamination was confirmed by immunocytochemical staining of rAAV/Ad co-infected 293 cells with anti-Rep antibodies. These studies, which had a sensitivity of 1 wtAAV in 10^{10} rAAV particles, demonstrated an absence of Rep staining as compared to pRep/Cap plasmid transfected controls.

Isolation and Structural Evaluation of AAV Circular Intermediates From HeLa Cells.

HeLa cells were grown in 35 mm dishes in DMEM media supplemented with 10% fetal calf serum (FCS). Cells were infected in the presence of 2% FCS at 80% confluency with recombinant AV.GFP3ori (MOI=1000 particles/cell, 1 x

10⁹ total particles/plate) and Hirt DNAs isolated as described by Duan et al. (1997) at 6, 12, 24, 48, and 72 hours post-infection. In experiments analyzing the effects of adenovirus, plates were co-infected with Ad.CMVLacZ (MOI=5000 particles/cell) in the presence of 2%FCS/DMEM. Zero hour

5 controls were generated by mixing 10⁹ particles of AV.GFP3ori with cell lysates prior to Hirt DNA preparation. Hirt DNA isolated at each time point was used to transform *E. coli* SURE cells (Stratagene, La Jolla, CA.). Typically, 1/10 of the Hirt DNA preparation was used to transform 40 ml of competent bacteria by electroporation. The resultant total number of bacterial colonies was quantified

10 for each time point and the structure of circular intermediates was evaluated for greater than 20 plasmid clones for each time point from two independent experiments. Structural determinations were based on restriction enzyme analysis using PstI, SphI, AseI single and double digests together with Southern blotting against GFP, stuffer, and ITR probes.

15 Evaluation of E2a and GFP gene expression in Hela cells.

E2a gene expression was evaluated by immunofluorescent staining of Hela cells superinfected with E1-deleted Ad.CMVLacZ (MOI= 0, 500, 5000 particles/cell). Briefly, cells were fixed in methanol at -20°C for 10 minutes followed by air drying. Cells were then incubated at room temperature with

20 hybridoma supernatant against Ad5 72kd DBP (Reich et al., 1983), followed by goat anti-mouse-FITC antibody (5 mg/ml) for 30 minutes at room temperature. In studies evaluating augmentation of AAV GFP transgene expression by adenovirus, Hela cells were harvested at 24 or 72 hours post-infection by trypsinization, resuspended in 2%FCS/PBS and evaluated by FACS analyses.

25 Thresholds were set using uninfected controls and the percentage and/or the average relative fluorescent intensity was determined by sorting greater than 10⁵ cells per experiment condition.

Sequence Analysis of AAV Circular Intermediates.

Sequence analysis of the ITR array within circular intermediates was

30 performed using primers EL118 (5'-CGGGGGTCGTTGGGCGGTCA-3'; SEQ ID NO:1) and EL230 (5'-GGGCGGAGCCTATGGAAAA-3'; SEQ ID NO:2) which are nested to 5' and 3' ITR sequences, respectively. Both circular and linearized (with SmaI which cuts within ITR sequences) plasmids were

sequenced.

Results

Construction of rAAV Shuttle Vector and Isolation of Circular Intermediates.

To circumvent the inability to retrieve pre-integration intermediates or as
5 stable episomal forms resistant to nuclease digestion, an alternative strategy was
developed to “trap” circular intermediates using a recombinant AAV shuttle
vector. Recombinant AV.GFP3ori virus (Figure 1B) was generated from a
cis-acting plasmid (pCisAV.GFP3ori, Figure 1A) by co-transfection in 293 cells
with *trans*-acting plasmids encoding Rep and Cap viral genes. This viral vector
10 (AV.GFP3ori) encoded the green fluorescent protein (GFP) reporter gene, a
bacterial origin of replication (*ori*), and the bacterial ampicillin-resistance gene.
Ori and ampicillin-resistance sequences encoded in this virus allow for the
rescue of circular AAV genomes formed during the transduction process.

To test this strategy, Hela cells were infected with AV.GFP3ori
15 (MOI=1000 particles/cell) and the abundance of circular intermediates was
evaluated following transformation of low molecular weight cellular Hirt DNA
into *E. coli* SURE cells. The presence of circular intermediates was inferred by
retrievable ampicillin-resistant bacterial colonies. Structural features of circular
intermediates were determined by restriction enzyme analysis and Southern
20 blotting with various regions of the provirus, including GFP, Stuffer, and ITR
sequences.

The predominant circular form isolated after transduction of Hela cells
with AV.GFP3ori consisted of 4.7 kb monomer-sized molecules (Figure 1C).
SphI digestions of these circular intermediates yielded characteristic 300 bp
25 bands which hybridized to an ITR probe on Southern blots (Figure 2A). PstI,
SphI, AseI single and double digests together with Southern blot analysis using
GFP, Stuffer (data not shown), and ITR (Figure 2A) probes confirmed the
structure of the circular intermediates as head-to-tail monomer genomes (Figure
1C). In particular, PstI digests together with ITR Southern blots distinguish
30 these head-to-tail circular intermediates from head-to-head or tail-to-tail circular
dimers. Similar results obtained with studies on AV.GFP3ori infected 293 cells
and primary fibroblasts have confirmed that monomer head-to-tail circular
intermediates were also the most abundant form in these cell types.

Because the predicted molecular weight of an intact head-to-tail ITR SphI fragment would be approximately 360 bp, an anomalous migration in agarose gels might be due to the high secondary structure of inverted repeats within ITRs. To this end, the head-to-tail orientation of the ITRs, as predicted
5 by Southern blot analysis, was confirmed using several sequencing strategies. First, the SphI ITR hybridizing fragment of a circular intermediates was subcloned into a secondary plasmid vector and sequenced with primers outside the ITR cloned sequences. These findings confirmed the head-to-tail orientation of ITRs. Additionally, sequence was obtained directly from six monomer
10 circular intermediate clones using primers internal to both the 5' and 3' ITRs (Figure 2C). In these studies, circular intermediates were digested with SmaI and the linear 4.6 kb plasmid was gel isolated prior to sequencing. SmaI digestion (which relaxed the secondary structure of ITRs) was necessary to obtain sequence information within the ITRs. The sequencing results presented
15 in Figure 2C confirmed the orientation of head-to-tail ITR arrays in these intermediates.

Interestingly, sequencing also revealed several consistent base pair (bp) changes in four of the six clones analyzed (Figure 2C). These four clones (p79, p81, p87, and p88) had consistent two bp changes within the D-sequence [G->A
20 (122 bp) and A->G (125 bp)], which always occurred together with the bp alterations in the p5 promoter [A->G (114 bp) and A->C (115 bp)]. No other consistent bp changes were noted except for two clones (p79 and p88) which demonstrated mutations just outside the 3' ITR D-sequence [T->G (381 bp) and T->C (383 bp)].

25 Although head-to-tail circular intermediates were the most abundant forms present in Hirt DNA from rAAV infected Hela cells, several less frequent structures were also detected. These included monomer circularized AAV genomes with one (p190) and three ITRs (p345) arranged in a head-to-tail fashion as well as several clones with an unknown structure lacking complete
30 ITRs (p340) (Figure 2A). Such diversity within the ITR array may represent homologous recombination *in vivo* or in bacteria during amplification. However, previous studies demonstrating similar variations in ITR sequences of head-to-tail integrated genomes, suggest that such changes in the length of the

ITR array may occur *in vivo* (Duan et al., 1997) Additionally, less frequent head-to-tail circularized multimer forms were predicted based on the variation in migration patterns of uncut plasmids which gave identical restriction patterns.

Results shown in Figure 2B confirmed the existence of monomer and dimer head-to-tail circular intermediates using partial digestion with an enzyme which cuts once in the AAV genome (AseI). Cumulative analysis of greater than 200 independently isolated circular intermediates from Hela cells demonstrated that head-to-tail circular AAV genomes occurred in greatest abundance as monomers (92%) and less frequently as multimers of greater than one genome (8%).

To establish that head-to-tail circular intermediates were formed *in vivo* and not by non-specific bacterial recombination of linear AAV genomes present in the Hirt DNA, a set of reconstitution experiments was performed by which the same number of rAAV particles used for infection experiments were spiked into Hela cell lysates prior to Hirt preparations. In these studies, background bacterial amplification of Hirt DNA spiked with linear rAAV genomes was negligible (Figure 3D) and of the few isolated colonies obtained from these controls, none had a predicted head-to-tail structure as assessed by Southern blot restriction enzyme analysis (Figure 3E). Additionally, reconstitution experiments transforming bacterial with linearized dsDNA AAV genomes did not give rise to significant levels of replication competent plasmids or the characteristic head-to-tail structure associated with AAV circular intermediates. These findings confirm that circular intermediates do not likely arise from non-specific recombination or ligation events with either ssDNA or dsDNA linear AAV genomes in bacteria. Additional control experiments, demonstrating the lack of stuffer hybridizing sequences in AAV circular intermediates by Southern blotting, also confirm that these structures do not arise from contamination of viral stocks with pCisAV.GFP3ori plasmid.

The formation of head-to-tail circular AAV intermediates is augmented by superinfection with E1-deleted adenovirus.

Many aspects of the wtAAV growth cycle are affected by helper adenovirus, including AAV DNA replication, transcription, splicing, translation, and virion assembly. Such studies have provided concrete evidence that a subset of Ad early gene products provide helper functions for the wtAAV lytic cycle,

including: E1a, E1b, E2a, E4 ORF6 and VA1 RNA (Muzyczka, 1992). In this regard, one of the most critical factors which is required for AAV replication is the 34 kD E4 protein (ORF6). Recent observations on the helper function of Ad in rAAV transduction have also demonstrated that Ad E4 ORF6 is essential for the augmentation of rAAV transgene expression seen with adenovirus co-infection (Ferrari et al., 1996; Fisher et al., 1996). According to these reports, the rate-limiting step enhanced by these adenoviral proteins is the conversion of single stranded AAV genomes to double stranded forms.

Studies evaluating the kinetics of rAAV circular intermediate formation demonstrated a time-dependent increase in abundance which peaked at 24 hours post-infection in Hela cells and coincided with the onset of GFP transgene expression (Figure 3). To better understand the cellular mechanisms associated with AAV circular intermediate formation, the effects of adenoviral co-infection on this process were evaluated. The extent of transgene expression and circular intermediate formation in AV.GFP3ori infected Hela cells with or without co-infection with E1-deleted recombinant adenovirus was compared.

Although E1-deleted adenoviruses are severely handicapped in their ability to synthesize viral gene products, at high MOIs of >5000 significant E2a protein expression was noted (Figure 3A). As an indicator of transgene expression, the abundance and average relative intensity of GFP positive cells was determined against mock infected controls by fluorescent microscopy (Figure 3B) and FACS analysis (Figure 3C) at 72 hours post-infection. In accord with previous reports demonstrating augmentation in rAAV transgene expression by adenovirus (Ferrari et al., 1996; Fisher et al., 1996), the extent of GFP transgene expression was dramatically increased at doses of adenovirus which led to viral gene expression (MOI>5000; Figures 3A-C). Additionally, persistence of rAAV transgene expression was also augmented by co-infection with E1-deleted adenovirus, as determined by GFP-expressing colony formation following serial passages (Figure 3C).

If circular intermediates represent a molecular form of rAAV important for efficient and/or persistent transgene expression, augmentation of rAAV transgene expression by adenovirus might also modulate circular intermediate formation. In these studies, the abundance and time course of AAV circular

intermediate formation was evaluated following superinfection with Ad.CMVLacZ. Results from these experiments are shown in Figure 3D, which represents the total number of bacterial colonies (per 35 mm plate) obtained following transformation of *E. coli* with Hirt DNA isolated from Hela cells infected with AV.GFP3ori (1000 DNA particles/cell) with or without co-infection with Ad.CMVLacZ (5,000 particles/cell). An MOI of 5000 Ad particles/cell was chosen for these experiments since this level of adenovirus led to minimal cytopathic effect (CPE) with high levels of E2a expression.

These studies demonstrated a nearly 2-fold augmentation by Ad.CMVLacZ in the total abundance of AAV rescued plasmid intermediates in *E. coli* (Figure 3D). Southern blot restriction enzyme analysis demonstrated that the predominant forms in both the presence and absence of adenovirus were head-to-tail monomer circular intermediates containing the diagnostic 300 bp ITR fragment following SphI digestion (Figure 3E). Additionally, results demonstrated that adenovirus co-infection led to an earlier time of onset and increased stability of AAV head-to-tail monomer circular intermediates (Figures 3E and F). For example, at 6 hours post-infection, head-to-tail circular intermediates were only present in Hela cells co-infected with adenovirus. Furthermore, a decline in the percentage of head-to-tail circular intermediate clones was seen at 48-72 hours post-AAV infection in the absence of adenovirus. In contrast, this decline was significantly blunted by the presence of helper adenovirus (Figure 3F). Based on these findings, it was concluded that certain adenoviral proteins produced by superinfection with E1-deleted adenovirus were capable of modulating circular intermediates formation and stability during rAAV transduction.

Discussion

In the present study, it was shown that circularization of linear AAV genomes occurs during rAAV transduction. Circularization appears to predominately occur as head-to-tail monomer genomes. However, the existence of less abundant circular multimer forms suggests that recombinational events subsequent to the initial infection may drive concatamerization of circular genomes. The diversity in the length of ITR arrays found within circular intermediates (i.e., 1-3 ITRs) also supports the notion that these forms may be

highly recombinagenic. Of mechanistic interest in the formation of circular intermediates is the uniformity of mutations observed in the D-sequences and nearby p5 promoter region and the confinement of these mutations to the 5'-ITRs. Although the etiology of these base pair changes is unknown, their uniformity suggests that they may have a direct role in the formation of circular intermediates and in increased stability. Recent findings, which suggest that an endogenous host single strand D-sequence binding protein is important in rAAV transduction, lend support to the potential involvement of this sequence in circular intermediate formation (Wang et al., 1997; Qing et al., 1998). Furthermore, it remains to be determined whether the *in vivo* formation of AAV circular intermediates occurs through the circularization of single or double stranded AAV genomes.

By analogy, retroviral transduction intermediates have striking similarities to the current findings with AAV. Three DNA forms have been isolated following retroviral infection, including linear DNA with long terminal repeats (LTRs) at both ends, circular DNA with one LTR, and circular DNA with multiple LTRs (Panganiban, 1985). Although it is disputed which of these forms are the direct precursor to integration, the existence of circular retroviral genomes which also have similar repeat regions at the ends of their genomes suggests the potential for common mechanisms with the formation of AAV circular intermediates. These AAV circular intermediates could act as integration precursors and/or stable episomal genomes.

The head-to-tail ITR structures found in AAV circular intermediates are most characteristic of latent integrated AAV genomes. In contrast, lytic phases of AAV growth are typically associated with head-to-head and tail-to-tail replication form genomes. Hence, it is likely that circular intermediates represent a latent aspect of the AAV life cycle. The finding that co-infection with adenovirus leads to increased abundance and stability of AAV circular intermediates suggests a novel link between adenoviral helper functions and latent infection of AAV.

Aspects of inverted head-to-tail ITRs, which include palindromic hairpins similar in structure to "Holliday-like" junctions, might impart recombinagenic activity which aids in viral integration. Such Holliday junctions

have been shown to play critical roles in directing homologous recombination in bacteria through the processing of recombination intermediates by RuvABC proteins (West, 1997; Lee et al., 1998). Interestingly, a mammalian endonuclease, analogous to bacterial RuvC resolvase, has also been isolated from cell lines (Hyde et al., 1994). Despite the theoretical considerations which might suggest that circular AAV genomes have characteristics of preintegration intermediates, a study with recombinant retrovirus has demonstrated that palindromic LTR-LTR junctions of MMLV are not efficient substrates for proviral integration (Lobel et al., 1989). Nonetheless, circular AAV genomes have been previously proposed as integration intermediates based on proviral structure (Linden et al., 1996).

Example 2

Methods

15 Production of rAAV Shuttle Vector.

The *cis*-acting plasmid (pCisAV.GFP3ori) used for rAAV production was generated by subcloning the Bsp1201/Not I fragment (743 bp) of the GFP transgene from pEGFP-1 (Clontech) between the CMV enhancer/promoter and SV40polyA by blunt-end ligation. A 2.5 kb cassette containing beta-lactamase and bacterial replication origin from pUC19 was blunt ligated down-stream of GFP reporter cassette. The ITR elements were derived from pSub201.2. The entire plasmid contains a 4.7 kb AAV component flanked by a 2 kb stuffer sequence. The integrity of ITR sequences was confirmed by restriction analysis with SmaI and PvuII, and by direct sequencing using a modified di-deoxy procedure which allowed for complete sequence through both 5' and 3' ITRs. Recombinant AAV stocks were generated by co-transfection of pCisAV.GFP3ori and pRep/Cap together with co-infection of recombinant Ad.CMVlacZ in 293 cells. The rAV.GFP3ori virus was subsequently purified through 3 rounds of CsCl banding as described in Duan et al., 1997. The typical yields from this viral preparation were 10¹² DNA molecules/ml.

DNA titers were determined by viral DNA slot blot hybridization against GFP ³²P-labeled probe with copy number plasmid standards. The absence of helper adenovirus was confirmed by histochemical staining of rAAV infected

293 cells for beta-galactosidase, and no recombinant adenovirus was found in 10^{10} particles of purified rAAV stocks. The absence of significant wtAAV contamination was confirmed by immunocytochemical staining of rAAV/Ad co-infected 293 cells with anti-Rep antibodies. Transfection with pRep/Cap was used to confirm the specificity of immunocytochemical staining. No immunoreactive Rep staining was observed in 293 cells infected with 10^{10} rAAV particles.

Isolation of AAV Circular Intermediates From Muscle.

The tibialis anterior muscle of 4-5 week old C57BL/6 mice were infected with AV.GFP3ori (3×10^{10} particles) in Hepes buffered saline (30 μ l). GFP expression was analyzed by direct immunofluorescence of freshly excised tissues and/or in formalin-fixed cryopreserved tissue sections in four independently injected muscles harvested at 0, 5, 10, 16, 22 and 80 days post-infection. Tissue sections were counter-stained with propidium iodide to identify nuclear DNA. Hirt DNA (Hirt, 1967) (20-ml per muscle sample) was isolated from at least three independent muscle specimen for each time point and used to transform *E. coli* SURE cells using 3 ml of Hirt with 40-ml of electrocompetent bacterial (approximately 1×10^9 cfu/ μ g DNA, Stratagene Inc.). The resultant total number of bacterial colonies was quantified for each time point and the abundance of head-to-tail circular intermediates was evaluated for each time point (> 20 bacterial clones analyzed) by PstI, AseI, SphI, and PstI/AseI digestion, and confirmed by Southern blot analysis using ITR, GFP and stuffer probes. The head-to-tail configuration in typical clones were also confirmed by dideoxy sequencing using primers EL118 (5'-CGGGGGTCGTTGGGCGGTCA-3'; SEQ ID NO:1) and EL230 (5'-GGGCGGAGCCTATGGAAAA-3'; SEQ ID NO:2) which are nested to 5' and 3' ITR sequences, respectively. Zero hour controls were generated by mixing 3×10^{10} particles of AV.GFP3ori with control uninfected muscle lysates prior to Hirt DNA preparation. As described in Table 1, a number of additional controls for were performed to rule out non-specific recombination of linear AAV genomes in bacteria as a source for isolated circular intermediates.

Table 1. Control Experiments for Rescue of Circular Intermediates in Bacteria

	Type of Input DNA	Source of DNA	Number of Molecules	Number of Amp Resistant Bacterial Colonies	Presence of Head-to-Tail Circular Intermediates
5	Purified rAVV	Hirt from Infected Muscle (22 day)	3×10^{10}	approximately 5×10^3	Yes
	Purified rAAV	Virus reconstituted into Uninfected Muscle Hirt ^a	3×10^{10}	0	No
10	Linear ssDNA Encompassing rAAV Genome ^b	Isolated from Purified Virus	3×10^{10}	2	No
	Linear dsDNA Encompassing Entire rAAV Genome	Isolated from proviral plasmid (HindIII/PvuII) ^c	3×10^{10}	3	No
15	Linear dsDNA Encompassing Entire rAAV Genome + ligase ^d	Isolated from proviral plasmid (HindIII/PvuII)	3×10^{10}	$>6 \times 10^3$	Yes

20.

^a Purified virus was reconstituted into muscle homogenates prior to preparation of Hirt DNA.

25 ^b Viral DNA predominantly contained single stranded genomes as evident by Southern blot analysis against with ITR probe. However, small amount of dsDNA AAV genomes also existed and are likely due to reannealing of single stranded genomes during preparation. Purified viral DNA concentrations were determined by OD₂₆₀ and 75 ng representing approximately 3×10^{10} viral genomes were used for transformation of bacteria.

30

35 ^c HindIII/PvuII digestion was used to remove the entire rAAV genome from pcisAV.GFP3ori. HindIII and PvuII leave 10 and 0 bps of flanking sequence outside the 5' and 3' ITRs, respectively. The linear dsDNA fragment (4.7 kb) was gel isolated following blunting with T4 DNA polymerase and the DNA concentration determined by OD₂₆₀. One hundred and fifty ng of linear fragment representing approximately 3×10^{10} viral genomes were used for transformation of bacteria.

^d Linear dsDNA viral genomes (HindIII/PvuII blunted fragment) were treated with T4 DNA ligase prior to transformation of bacteria.

^e The presence of head-to-tail circular AAV intermediates were confirmed by restriction enzyme digestion (AseI, PstI, and SphI) and Southern blotting against ITR probe.

Fractionation of muscle Hirt DNA preparations.

Preparative-scale fractionation of the muscle Hirt DNA was performed by 1% agarose gel electrophoresis using the Bio-Rad Mini Prep Cell (Catalog #170-2908). A 4.5 ml (10.5 cm) tubular gel containing 1 x TBE buffer was poured according to manufacturer's specification. A total of 20 ml Hirt preparation from one entire muscle sample was loaded on top of the gel. Electrophoresis was carried out at a constant current of 10 mA over a period of 5 hours. Sample eluent was drawn from the preparative gel apparatus by a peristaltic pump at a rate of 100 ml/min and eluted into a fraction collector at 250 ml/fraction. The collected DNA was subsequently concentrated by standard ethanol precipitation and used to transform SURE bacterial cells by electroporation as described above.

In vitro Persistence of AAV Circular Intermediates.

Transgene expression and persistence of AAV circular intermediate plasmid clones were evaluated following transient transfection in HeLa and 293 cells. Subconfluent monolayers of HeLa cells in 24-well dishes were transfected with 0.5 mg of either AAV circular intermediates (p81 or p87) or pCMVGFP using Lipofectamine (Gibco BRL Inc.). The cultures were then incubated for 5 hours in serum free DMEM followed by incubation in DMEM supplemented with 10% fetal bovine serum. All plasmid DNA samples used for transfections were spiked with pRSVlacZ (0.5 mg) as an internal control for transfection efficiency. At 48 hours post-transfection, cells were passaged at a 1:10 dilution and allowed to grow to confluency (day 5), at which time GFP clones were quantified for size and abundance using direct fluorescent microscopy. The percent of beta-galactosidase-expressing cells was also quantified at this time point by X-gal staining. At 5 days, cells were passaged an additional time (1:15 dilution) GFP clones were quantified again at day 10. The persistence of

plasmid DNA at passage-5, 7, and 10 days post-transfection was evaluated by Southern blot analysis of total cellular DNA using ^{32}P -labeled GFP probes. To determine whether the head-to-tail ITR array within circular intermediates was responsible for increases in the persistence of GFP expression, the head-to-tail ITR DNA element was subcloned into the pGL3 luciferase plasmid to generate pGL3(ITR). The head-to-tail ITR DNA element was isolated from a monomer circular intermediate (p81) by AatII and HaeII double digestion and subsequently inserted into the SalI site of pGL3 (Promega) by blunt ligation. The resultant plasmid pGL3(ITR) contains the luciferase reporter and head-to-tail ITR element 3' to the polyA site. The integrity of the ITR DNA element within this plasmid was confirmed by sequencing. The persistence of transgene expression from pGL3(ITR) was compared to that of pGL3 by luciferase assays on transiently transfected Hela cells as described above and analyzed at 10 days (passage-2). Transfection efficiencies were normalized using a dual renilla luciferase reporter vector (pRLSV40, Promega).

Results

AAV Circular Intermediates Represent Stable Episomal Forms of Viral DNA Associated with Long-term Persistence of Transgene Expression in Muscle.

To evaluate the molecular characteristics of rAAV genomes in muscle, a rAAV shuttle viral vector (AV.GFP3ori) was utilized which harbors an ampicillin resistance gene, bacterial origin of replication, and GFP reporter gene (Figure 1A). This recombinant virus was used to evaluate the presence of circular intermediates by bacterial rescue of replication competent plasmids. In these studies, delivery of AV.GFP3ori (3×10^{10} particles) to the tibialis muscle of mice led to GFP transgene expression which peaked at 22 days and remained stable for at least 80 days (Figure 4A). These results confirmed previous successes in rAAV mediated gene transfer to muscle (Kessler et al., 1996; Herzog et al., 1997; Xiao et al., 1996; Clark et al., 1997; Fisher et al., 1997). The formation of circular intermediates was evaluated by *E. coli* transformation of Hirt DNA harvested from muscle at 0, 5, 10, 16, 22, and 80 days post-infection with AV.GFP3ori.

In these muscle samples, circular intermediates were found to have a characteristic head-to-tail structure with 1-2 ITR repeats. The most abundant

form included two inverted ITRs within a circularized genome (Figure 4B, clone p17). This figure also depicts a less frequent form (< 5%) of circular intermediates observed, p439, with undetermined structure. When this type of replication competent plasmid was seen, it was not included in the

5 quantification of head-to-tail circular intermediates since its structure could not be conclusively determined. The total abundance of muscle Hirt derived head-to-tail circular intermediates (with 1-2 ITRs) demonstrated a time-dependent increase that peaked with transgene expression at 22 days and slightly decreased by day 80 (Figure 5A). Increased diversity in the length of

10 ITR arrays within circular intermediates was seen at longer time points. For example, Figure 5B demonstrates several isolated circular intermediates with 1-3 ITRs isolated from 80 days muscle Hirt samples. This is in contrast to the more uniform structure of circular intermediates with two ITRs in a head-to-tail conformation at 5-22 days post-infection.

15 To evaluate the potential for artifactual rescue of linear rAAV genomes by recombination in bacteria, several control experiments were performed. First, uninfected control muscle Hirt preparations, spiked with an equal amount of rAAV virus used for *in vivo* infection of muscles, failed to give rise to replicating plasmids following transformation of *E. coli*. Second, when a blunted

20 linear double stranded HindIII/PvuII fragment isolated from pcisAV.GFP3ori (encompassing the entire rAAV genome) was used to transform bacteria, no ampicillin resistant bacterial colonies were obtained. The addition of T4 ligase to this fragment, however, led to significant numbers of bacterial colonies. Third, when purified single stranded rAAV DNA was used for transformation,

25 no bacterial colonies were obtained. As summarized in Table 1, these results confirm that in the absence of productive infection, rAAV genomes themselves are incapable of recombining into replication competent plasmids in bacteria. Hence, *in vivo* circularization of rAAV genomes is a prerequisite for rescuing autonomously replicating plasmids in *E. coli* with this shuttle vector.

Molecular weight of circular intermediates suggest a conversion from monomer to multimer forms over time.

To further characterize the circular intermediates isolated from muscles, Hirt samples from 22 days and 80 days post-infected muscles were size fractionated by continuous-flow gel electrophoresis (BioRad). As shown in Figure 6, the majority of circular intermediates at 22 days post-infection size fractionated at a molecular weight of less than 3 Kbp. Very few clones were isolated from fractions between 3 to 5 kb and no clones were obtained from fractions larger than 5 kb at this time point. Furthermore, this size fractionated molecular weight of *in vivo* Hirt derived circular intermediates at 22 day time points correlated with that of head-to-tail monomer undigested circular intermediate plasmids rescued in bacteria from this same time point (approximately 2.5 kb). These data suggest that at early time points post-infection in muscle, the predominant form of circular intermediates likely occurs as monomer genomes. The lower mobility of this fraction as compared to replication form monomer (Rfm=4.7 kb) and dimer (Rfd=9.4 kb) genomes provides indirect evidence that these forms are not responsible for rescued plasmids in these Hirt samples. Interestingly, when 80 day muscle Hirt samples were size fractionated, more clones were retrieved from higher molecular weight fractions ranging from 3-12 kb (Figure 6). This shift in the molecular weight of circular intermediates indicates the potential for recombination between monomer forms in the generation of large circular multimer genomes. Such concatamerization has been previously observed in muscle and has traditionally been hypothesized to involve linear integrated forms of the AAV genome (Herzog et al., 1997; Xiao et al., 1996; Clark et al., 1997; Fisher et al., 1997). This data sheds new light on the molecular characteristics of these persistent AAV genomes and suggests that they are in fact circular and episomal. Based on yields of retrievable circular plasmids reconstituted in Hirt DNA, the efficiency of bacterial transformation, and the initial inoculum of virus, we estimate that approximately 1 in 400 viral DNA particles circularize following infection in muscle (Table 2).

Table 2. Yield of Circular Intermediate Isolation from Hirt DNA

5	Bacterial Transformation	Starting Number of Plasmid or AAV Genomes	Actual Number of Amp ^r cfu	Adjusted Yield
	Hirt DNA from rAAV Infected Muscle ^a	3 x 10 ¹⁰ molecules	5 x 10 ³ cfu	5 x 10 ⁵ cfu ^e
	Hirt DNA + 230 ng LacZ Plasmid ^{b, c}	3 x 10 ¹⁰ molecules	2 x 10 ⁶ cfu ^d	2 x 10 ⁸ cfu
10	230 ng LacZ Plasmid ^c	3 x 10 ¹⁰ molecules	2 x 10 ⁸ cfu	-----

^a The actual amount of Hirt used for transformation was 3/20 the entire Hirt DNA. The numbers have been adjusted to reflect viral inoculum and yields for the entire muscle.

15

^b Plasmid DNA was spiked into mock infected muscle homogenates prior to isolation of Hirt DNA. This reconstituted Hirt DNA was then used for transformation of bacteria.

20

^c The actual microgram amounts of plasmid used in reconstitution experiments was 10 ng. The numbers have been adjusted for comparison to normalize the number of plasmids genomes to that used in AAV experiments. Control LacZ plasmid was approximately 7000 bp with a molecular weight of 4.6 x 10⁶ g/mole.

25

^d The average of several experiments indicates an approximate 100-fold reduction in the number of cfu recovered from bacterial transformations with DNA isolated from Hirt extract spiked with plasmids as compared to transformation with an equivalent amount of plasmid DNA alone.

30

^e Adjusted yield indicate approximately 1 in 400 AAV genomes circularize *in vivo*.

35

Given the fact that not all rAAV particles likely contain functional DNA molecules and intermediates may integrate, these calculations may represent an underestimation.

AAV Circular Intermediates Demonstrate Increased Persistence as Plasmid Based Vectors.

40

Based on the finding that circular AAV intermediates were associated with long term persistence of transgene expression in muscle, rAAV circular

head-to-tail intermediates may be molecular structures of the AAV genome associated with the latent life cycle and increased episomal stability. Several aspects of the structure of AAV circular intermediates may account for their increased stability *in vivo*. First, circularization of AAV genomes may create a
5 nuclease resistant conformation. Secondly, since the only viral sequences contained within circular intermediates are the head-to-tail ITR array, these sequences might bind cellular factors capable of stabilizing these structures *in vivo*. Several studies have demonstrated increased persistence of transgene expression with plasmid DNA encoding viral ITRs (Philip et al., 1994; Vieweg
10 et al., 1995). The results described above provide a functional explanation for the increased persistence through the association with circular intermediate formation as part of the AAV life cycle.

To more closely evaluate the persistence of AAV head-to-tail circular intermediates, several *in vitro* experiments were performed by transfecting these
15 intermediates into Hela cells and assessing the stability of plasmid DNA and transgene expression by GFP clonal expansion. Results from Hela cell transfection experiments demonstrated that two monomer head-to-tail circular intermediates (p81 and p87) studied gave rise to a 10-fold higher number of five and ten day transgene-expressing clones, as compared to a control pCMVGFP
20 plasmid lacking the ITR sequences (Figures 7A and B). Additionally, the size of GFP positive colonies at 5 days post-transfection was three-fold larger in Hela cells transfected with p81 and p87, as compared to the pCMVGFP control vector (Figures 7A and B). These studies suggest the AAV circular intermediates have increased stability of transgene expression and substantiate findings in muscle.

25 To confirm the increased molecular persistence of head-to-tail circular intermediates following transfection into Hela cells, total DNA (low and high molecular weight) was isolated from cultures of pCMVGFP and p81 transfected Hela cells at various passages post-transfection and analyzed by Southern blotting. Southern blots hybridized to ³²P-labeled GFP probes demonstrated a
30 significantly higher level of p81 plasmid DNA at passage-7 as compared to the control vector lacking the head-to-tail ITR sequence (Figure 7C). The majority of signal in undigested DNA samples was associated with a 4.7 kb band migrating at the approximate size of the uncut monomer plasmids. Together

with the fact that the majority of signal from all cell cultures in Figure 7C disappeared by passage-10, these data suggest that these plasmids predominantly remained episomal. Thus, in both muscle and Hela cells, increased persistence of AAV circular intermediates is correlated with stable transgene expression.

5 ITR arrays are responsible for increased persistence.

To investigate whether the head-to-tail ITR DNA element was responsible for the increased persistence of circular intermediates, we cloned this DNA element into a secondary luciferase vector (pGL3) to give rise to pGL3(ITR). Transient transfection experiments in Hela cells demonstrated a
10 five-fold increase in the persistence of luciferase expression in serially-passaged cultures at 10 days in pGL3(ITR) as compared to that of pGL3 transfected (Figure 7D). These findings support the hypothesis that the head-to-tail ITR DNA element contained within circular intermediates is responsible for mediating the increased persistence of transgene expression and suggest a
15 mechanism by which these molecular intermediates may confer stability to AAV genomes *in vivo*. Furthermore, increases in the stability of transgene expression conferred by this element appear to be primarily context independent, since the head-to-tail ITR element was 3' to the luciferase gene in pGL3(ITR) and 5' to the GFP transgene in AAV circular intermediates.

20 Discussion

Characterization of integrated proviral structures in different cell lines has demonstrated head-to-tail genomes as the predominant structural forms for both wild type and recombinant AAV (McLaughlin et al., 1988; Cheung et al., 1980; Duan et al., 1997). This is in contrast to the head-to-head and tail-to-tail
25 structures observed in AAV replication intermediates (Rfm and Rfd). Both Rfm and Rfd configurations have also been demonstrated in rAAV infected cells and enhanced conversion of ssAAV genomes to double stranded Rfm and Rfd forms has been suggested as a mechanism for augmentation of rAAV transduction by adenovirus in cell lines (Ferrari et al., 1996; Fisher et al., 1996). However, it is
30 plausible that the mechanisms responsible for the formation of Rfm and Rfd molecules are different from pathways which lead to long-term transgene expression. In support of this hypothesis is a recent study evaluating augmentation of rAAV transgene expression by adenovirus in liver (Snyder et

al., 1997). These studies have demonstrated that co-infection of the liver with adenovirus and rAAV enhances short term transgene expression while long term expression was no different than rAAV alone. The exact mechanism for the formation of head-to-tail circular intermediates is not clear, however similar structures have been demonstrated to act as pre-integration intermediates for retrovirus (Varmus, 1982). In this regard, circularized retroviral genomes with one and two viral LTRs have been proposed. In addition, circular pre-integration intermediates have also been suggested by recent studies on wtAAV integration (Linden et al., 1996b). The demonstration that circular intermediates exist in rAAV infected muscle explains several features of latent phase infection with rAAV vectors including proviral structure and stable episomal persistence.

Previous studies have suggested that rAAV genomes delivered to muscle might persist as head-to-tail concatamers (Herzog et al., 1997; Clark et al., 1997; Fisher et al., 1997). However, it is currently unknown whether these concatamers exist as free episomes or as integrated proviruses in the host genome. The results described above, i.e., demonstrating prolonged persistence of head-to-tail circular intermediates at 80 days post-infection, suggest that a large percentage of rAAV genomes may remain episomal. The conversion of monomer circularized genomes to larger circularized multimers appears to be an aspect associated with long term persistence and likely represents recombinational events between monomer intermediates. Although the bacterial rescue strategy was not capable of satisfactorily addressing the size of multimers, our modified approach to size fractionating Hirt DNA prior to bacterial rescue of intermediates lends support to this hypothesis. Additional supportive evidence for increased recombination over time is the finding that greater variability in the length of ITR arrays was observed at longer time points post-infection. For example, at 5-22 days the majority of circular intermediates contained 2 ITRs in a head-to-tail fashion. This is in contrast to 80 day time points where the lengths of ITR arrays ranges from 1-3 ITRs. Such diversity of ITR arrays in muscle infected with AAV has been previously found using PCR approaches (Herzog et al., 1997; Fisher et al., 1997). In addition, the 30% decline in the abundance of circular intermediates in muscle between 22 and 80 days also supports a hypothesis that these molecular forms of AAV may represent pre-integration

complexes.

Given the fact that circular intermediates had long term persistence in muscle, certain structural features of these intermediates may affect episomal stability of DNA. Previous studies have noted increased persistence of transgene expression from plasmids encoding AAV ITRs (Philip et al., 1994; Vieweg et al., 1995). However, the physiologic significance of this finding has remained elusive. The present study, demonstrating the head-to-tail ITR arrays isolated from AAV circular intermediates can confer increased episomal persistence to plasmids following transfection in cell lines, gives a mechanistic framework for ITR effects on plasmid persistence. Furthermore, the correlation that AAV circular intermediates have increased persistence in cell lines *in vitro*, lends support to the hypothesis that these structures represent stable episomal forms following rAAV transduction in muscle. Stability of circular intermediates *in vivo* might be mediated by the binding of cellular factors to "Holliday-like" junctions in ITR arrays which stabilize or protect DNA from degradation.

rAAV has been shown to be an efficient vector for expressing transgenes in various tissues in addition to muscle, such as brain, retina, liver, lung, and hematopoietic cells (Snyder et al., 1997; Muzyczka, 1992; Kaplitt et al., 1994; Walsh et al., 1994; Halbert et al., 1997; Koeberl et al., 1997; Conrad et al., 1996; Bennett et al., 1997; Flannery et al., 1997). Despite these advances in the application of rAAV, the mechanisms of *in vivo* rAAV-mediated transduction and persistence of transgene expression still remain unclear. Such questions as to the molecular state of rAAV following *in vivo* delivery is highly relevant to the clinical application of this viral vector. For example, should rAAV primarily persist as an randomly integrated provirus, the potential for insertional mutagenesis could present a major theoretical obstacle in the use of this vector due to the potential for mutational oncogenesis. The demonstration that rAAV can persist as episomes suggest that random integration and associated risks of malignancy may not be a major concern for this viral vector system. Additionally, the molecular determinants of AAV circular intermediates associated with increased persistence in cell lines appear to be contained within the DNA elements encompassing the inverted ITRs. The isolation of this naturally occurring viral DNA element, which forms as part of the AAV life

cycle and acts to stabilize circular episomal DNA, may prove useful in increasing the efficacy of both viral and non-viral gene therapy vectors.

Example 3

5 Evidence for Increased Episomal Persistence of AAV Circular Intermediates in a Model for *in utero* Plasmid-Based Gene Therapy

Persistence of AAV circular intermediates were assessed by injection of plasmid DNA directly into the pronucleus of fertilized *Xenopus* oocytes. Twenty-five ng of the p81 isolate of AAV circular intermediates was injected at
10 the single cell stage of fertilized *Xenopus* oocytes. This plasmid was compared to the proviral plasmid pCisAV.GFP3ori, which contains two ITRs separated by stuffer sequence in an alternative confirmation to ITRs in p81. Figure 13 depicts the persistence of GFP plasmids as assessed by direct fluorescence of GFP. At this state of tadpole development, the fertilized oocyte has expanded from a
15 single cell to approximately 10^6 cells.

These studies confirm that AAV circular intermediates (p81) confer a higher level of stability in development *Xenopus* oocytes than plasmids containing similar transcriptional elements and ITR sequences in an alternative confirmation. Given that in the case of p81 injected oocytes, tadpoles are
20 completely fluorescent, the data suggests that some level of integration may have occurred.

Example 4

25 Liposome Mediated Transfer of Vectors of the Invention to the Airway and Muscle

Studies evaluating the mechanisms of recombinant adeno-associated virus (AAV) transduction have identified a novel molecular intermediate responsible for episomal persistence. This intermediate is characterized by a circularized AAV genome with head-to-tail ITR repeats. Circular intermediates
30 of rAAV were identified using a recombinant shuttle vector capable of propagating circularized viral genomes in bacteria. Pivotal experiments in cell lines demonstrate that the formation and persistence of these circular intermediates are augmented in the presence of helper adenovirus. These

findings suggest that cellular factors induced by adenoviral gene expression may modulate both the formation and/or persistence of AAV circular intermediates. Furthermore, studies in muscle have demonstrated that following rAAV infection, the formation and persistence of AAV circular intermediates correlates with the onset and maintenance (at 80 days) of transgene expression, respectively. Moreover, a 300 bp fragment encompassing the head-to-tail inverted ITR repeats found in AAV circular intermediates when cloned into heterologous expression plasmids can confer increased stability to those plasmids in HeLa cells. The structural aspects of AAV circular intermediates may lead to development of non-viral, plasmid based, gene transfer vectors with increased persistence of transgene expression.

To determine whether AAV circular intermediates which differ in length and/or sequence of the ITR array are more efficacious plasmid based vectors for liposome-mediated gene transfer to the airway and muscle, several distinct forms of AAV circular intermediates are evaluated as plasmid-based delivery systems in three model systems of the airway including: 1) *in vitro* polarized primary airway epithelial monolayers, 2) mouse lung, and 3) human bronchial xenografts. Persistence is evaluated at both the level of transgene expression (using GFP and luciferase reporters) and at the level of episomal and integrated transgene derived DNA. Studies are performed to assess whether integration can be specifically enhanced by co-transfection with Rep DNA or mRNA. These studies also evaluate both the extent of integration and site specificity to AVS1 sites in chromosome 19 of human model systems.

Gene therapy using plasmid-based delivery systems have encountered several obstacles to efficient transgene expression. These obstacles include transient expression of transgenes and rapid degradation of DNA. In contrast, viruses have developed efficient mechanisms for transducing cells and expressing encoded viral genes. The molecular characteristics of AAV circular intermediates which confer increased persistence of transgene expression include a DNA element encompassing the head-to-tail ITR. Based on the findings that circular intermediates have increased episomal persistence in muscle following rAAV transduction, these structures may also have increased persistence as plasmid-based vehicles to the airway. Interestingly, several naturally occurring

mutations which are found in approximately 50% of AAV circular intermediates affect the stability of the intermediate.

Several findings evaluating the efficiency of AAV circular intermediate formation from recombinant viral vectors have suggested that these structures are augmented in abundance by the presence of the E2a adenoviral gene product. These molecular structures may represent preintegration intermediates which, in the case of wild-type AAV, would efficiently integrate into the cellular genome by Rep facilitated mechanisms. However, in the case of recombinant AAV genomes (in the absence of Rep proteins), evidence suggests that these structures have increased episomal stability. To test whether exogenous addition of Rep and/or E2a can increase the efficacy of AAV circular intermediates by modulating their stability and/or integration, co-transfection methods with Rep encoding plasmids and mRNA are conducted. Additionally, exogenously supplied E2a DNA binding protein (DBP) may also enhance stability of AAV circular intermediates. Rep may increase the integration of circular intermediates while E2a may increase their episomal stability. Several observations including the association of E2a DBP with AAV genomes in the nucleus support a direct interaction between DBP and AAV circular intermediates. Furthermore, if DBP associates with AAV circular intermediates, its encoded nuclear localization sequence (NLS) may enhance nuclear sequestration of these plasmids in the nucleus. Alternatively, E2a may act to alter the persistence of AAV circular intermediates through the induction of cellular factors which interact with the ITR array.

Liposome mediated gene transfer to the airway has considerable advantages due to the low level of toxicity. However, limitations include transient low level expression in differentiated airway epithelia. Despite this apparent limitation, several laboratories have had considerable success with the use of cationic liposome-mediated gene transfer in several animal models including mouse and rat lung, and numerous laboratories have pursued clinical trials, which suggested that these vehicles may show promise for gene therapy of the cystic fibrosis (CF) lung. Thus, delivery of the present vectors in plasmid form via liposomes may be a safe and effective vehicle for gene transfer to the airway.

To assess whether AAV circular intermediates may also have increased persistence in airway epithelial cells as seen in Hela cells, several distinct forms of circular intermediates delivered by liposome-mediated transfection into primary airway epithelial cells, are evaluated. Based on the diversity of ITR repeat elements between various isolated circular intermediates (i.e., including 0, 1, 2, and 3 ITRs), circular intermediates isolated from later time points in muscle may have been naturally selected for increased stability *in vivo*. Hence, the structural consistencies between AAV circular intermediates are identified which give increased persistence as plasmid based vectors for gene transfer.

10 Circular intermediates containing the GFP reporter gene and 1, 2, and 3 ITRs are transfected into primary airway cultures and polarized epithelial cell monolayers using the cationic lipid GL-67 (Genzyme Inc.). DNA to lipid ratios are optimized using a luciferase reporter. Additionally, the addition of EGTA, or the use of calcium-free media, can increase the extent of gene transfer about 10- fold, and may be included to enhance gene transfer to polarized epithelial 15 monolayers. To evaluate persistence and expression of transgenes from circular intermediates, direct fluorescent microscopy and Southern blotting of both Hirt and genomic DNA with GFP P³²-labeled probes are utilized. Proliferating cultures of primary airway epithelial cells can be passaged up to 4 times during 20 this analysis. In contrast, polarized epithelial monolayers are evaluated at 1 week intervals for DNA persistence for up to 6 weeks. Since GFP transgene expression may be low and difficult to detect by direct fluorescence, GFP is quantitated by fluorometer of cell lysates.

25 Following AAV transduction, circular intermediates may form within cells and certain structures of these intermediates may persist by virtue of affinity for cellular factors which bind at ITR arrays. If this is true, then it may be possible to select for and isolate optimal circular intermediates with increased persistence in airway cells by batch screening of circular intermediates pools from rAAV infected airway epithelia.

30 Primary airway epithelia cell cultures are infected with AV. GFP3ori (MOIs of 1000 to 10,000 DNA part/cell) and low molecular Hirt DNA is prepared at 5-15 days post-infection. Hirt DNA containing circular intermediates from rAAV infected cells is used to then transfect primary airway

epithelial cells from which Hirt DNA is prepared at 5-15 days post-transfection. This second Hirt isolation is then used to isolate replication competent plasmids following transformation into bacteria. This selection process may give rise to those populations of circular intermediates with increased episomal persistence
5 in airway epithelial cells. Selected clones of circular intermediate plasmids isolated by this procedure are then tested individually for increased persistence following liposome mediated transfection. These studies are performed in a batch type screening in 24 well plates using two serial passages for persistence. Once plasmids having increased persistence are isolated, their structure and
10 sequence of ITR arrays are characterized. Since screening is performed on small-scale cultures, it may be necessary to implement semi-quantitative screening for DNA persistence within the first round of transfection using PCR methods. Candidate plasmids with a high level of increased persistence as compared to control plasmids which lack ITR sequences but contain the identical
15 promoter-reporter element, are evaluated on a larger scale transfection amenable to analysis by Southern blotting of total DNA.

To evaluate selected circular intermediate structures *in vivo*, two models including mouse lung and the human bronchial xenograft are employed. 10 wk BalbC mice are transfected with GL-67/DNA complexes at a ratio of 25 µg
20 plasmid/25 µg lipid in an iso-osmotic solution of Dextrose. At 1, 5, 10, 15, and 20 days post-transfection lungs of mice are harvested for immunofluorescent detection of GFP in formalin fixed sections and for quantitative fluorometry of tissue lysates. Southern blots are employed to evaluate the persistence of plasmids in Hirt and genomic DNA. In addition to evaluating the persistence of
25 selected circular intermediates which have the highest level of persistence with *in vitro* models, luciferase constructs are evaluated in which the ITR array has been cloned either 5' or 3' to the reporter gene. Furthermore, the use of luciferase reporters allows for more sensitive assessment of transgene activity in cell lysates.

30 Similarly, *in vivo* persistence of transfected circular intermediates and heterologous plasmids containing ITR arrays found within circular intermediates is evaluated in human bronchial xenografts.

Findings evaluating the effects of adenoviral co-infection on circular intermediate formation and persistence have suggested that E2a DBP leads to a 10-fold increase in the abundance of circular intermediates as compared to E2 deleted virus. Furthermore, studies with E1-deleted virus have demonstrated that the persistence of circular intermediates in Hela cells is increased at 72 hours post-infection. These studies suggest that E2a DBP may augment circular intermediate formation and/or increase the stability of these structures by an unknown mechanism. E2a DBP may interact directly with circularized genomes and/or induce cellular factors which interact with sequences in these AAV genomes. Since DBP encodes an NLS, this protein may act to shuttle circular intermediates to regions of nucleus that allow for increased stability of these structures. NLS sequences have been shown to cooperatively interact with nucleolar targeting sequences and hence we will also evaluate if subnuclear targeting is important in maintaining the increased stability of circular intermediates containing ITR arrays. Furthermore, it is currently unknown where circular intermediates form in the cell and it remains plausible that they may form in the cytoplasm or nucleus. Hence, if DBP associates directly with circular intermediates, it may act as an NLS for DNA to enter the nucleus as well.

Several *in vitro* reconstitution models are used to investigate the interaction of circular intermediates with DBP and their affect on *in vivo* persistence following DNA transfection in Hela cells. Furthermore, results evaluating the affects of various mutant adenoviral vectors on circular intermediate and Rfm/Rfd formation have suggested that these two types of intermediates occur by independent pathways indicative of latent and lytic infection, respectively. In the setting of wild type AAV, circular intermediates may be pre-integration complexes, which in the presence of Rep, efficiently integrate into the host genome. In contrast, in the absence of Rep, circular intermediates may accumulate episomally in rAAV infected cells. To this end, methods of supplementing Rep function may be capable of enhancing integration of plasmid based delivery of AAV circular intermediates. Thus, experiments in which co-transfection of circular intermediate plasmids with Rep expression plasmids or mRNA are conducted.

To investigate whether DBP can augment the stability of circular intermediates by increasing targeting to the nucleus, a HeLa cell line (gmDBP6) is utilized which encodes an inducible E2a gene under a dexamethasone responsible element. This cell line gives rise to high levels of DBP in nuclear
5 extracts by Western blot following treatment with dexamethasone. gmDBP6 cells (+/- DEX) are transfected with various AAV circular intermediate plasmids containing 0, 1, 2, and 3 ITRs and total cellular and nuclear plasmid content evaluated by subcellular fractionation using Southern blotting against GFP
10 probes. The time course of these studies is initially within the range of 12 hours to 4 days post-transfection. Transgene expression is evaluated by fluorometry (in cell lysates), and fluorescent microscopy (in viable cells), for GFP and luminescence for luciferase. HeLa cells have demonstrated that immediate increases in transgene expression from AAV GFP circular intermediates as compared to control GFP plasmids occur as early as 24 hours post-transfection.
15 Thus, certain cellular factors may facilitate an immediate accumulation of circular intermediates in the nucleus. DBP may invoke this increase by either direct interactions with ITR sequences or by the induction of cellular factors. To evaluate the potential for direct interactions between DBP and circular intermediates, various form of ITR arrays found within circular intermediates are
20 end-labeled with γ -ATP³² and evaluated for binding by electrophoretic mobility shift assays to nuclear extracts from gmDBP6 cells (+/- DEX). Supershifts, with DBP antibodies and competition experiments with cold ITR sequences and non-specific DNA, are used as controls for specific binding.

In a second model system aimed at evaluating the potential of DBP for
25 shuttling and/or sequestering of circular intermediates to the nucleus, microinjection experiments in oocytes are performed with 50 ng of plasmid DNA of circular intermediates with and without 50 ng of DBP mRNA. Experiments initially evaluate the time course of GFP transgene expression (+/- DBP cRNA) by direct fluorescent microscopy. If major differences are seen,
30 quantitative fluorometry of individual whole oocytes in 96 well plates is conducted. Similar studies on nuclear targeting in the presence of DBP can also be evaluated in this model by pooling microinjected oocytes for nuclear isolation and Southern blot analysis.

A third experimental model to evaluate nuclear targeting and/or accumulation of circular intermediate vectors in the presence and absence of DBP involves the microinjection of fluorescently labeled plasmid DNA into the cytoplasm and real time imaging to follow the nuclear accumulation of DNA.

- 5 The DNA fluorescent dye, TOTO-1, is used to label DNA prior to injection. This dye forms an extremely stable complex with negligible diffusion and re-incorporation into nuclear DNA following transfection into polarized airway epithelial cell monolayers. Co-localization of DBP with wtAAV DNA genomes at focal hot spots within the nucleus supports the observation that nucleolar
- 10 targeting may be important for persistence. These experiments are also performed in primary airway epithelial cells and *in vivo* models of the airway by either co-transfection of circular intermediates with DBP expressing plasmids and/or mRNA.

- The effects of Rep co-transfection on the integration of circular
- 15 intermediate plasmids is also evaluated. Two methods are used to express Rep including: 1) co-transfection with Rep expressing plasmids, and 2) co-transfection with Rep encoding mRNA. Initially, Hela, CFT1, and IB-3 cells are tested, as transformed cells may be more amenable to expansion and evaluation of integration. Both CFT1 and IB-3 cells represent airway epithelial cells.
- 20 Experiments are performed by cationic liposome (GL-67) mediated transfection of circular intermediate DNA with varying doses of a Rep-containing expression vector, e.g., pCMVRep. The extent of integration is also evaluated by two criteria, Southern blotting of Hirt and genomic DNA and clonal expansion of GFP expressing cells. Since Southern blot has an approximate limit of
- 25 sensitivity of 1 integrated plasmid molecule per 10 cellular genomes, clonal expansion may be necessary to evaluate persistence in less transfectable cells such as CFT1 and IB-3 cells. Cell lines are evaluated over the course of 1-10 passages.

- Sustained expression of Rep by plasmid mediated co-transfection may be
- 30 toxic to cells, hence co-transfection with Rep mRNA is also evaluated. Cationic liposome:mRNA mediated transfection has been previously shown to work in cell lines and although the level of expression is much more transient than for DNA, in these studies it may be an advantage. Initial studies are performed with

in vitro transcribed Rep mRNA alone to evaluate the μ g amount of mRNA needed for Rep expression as determined by Western blot. Once the threshold for detectable Rep expression is established, increasing amounts of Rep mRNA are co-transfected with circular intermediate DNA. Similar assays are used as described above to evaluate the extent of AAV circular intermediate integration. If findings suggest that increased integration is facilitated by Rep, the site specificity of this integration can be evaluated by cloning GFP expressing cells after the 10th passage by serial dilution. These GFP expressing clones are expanded and genomic Southern blots assessed with both GFP and AVS1 specific probes. By evaluating a number of restriction enzymes which either do not cut or cut once within the circular intermediate plasmid, it will be determined whether integration has occurred at the AVS1 loci.

To test whether secondary structure rather than primary sequence is the important determinant of increased episomal stability of AAV circular intermediates, synthetic DNA sequences are generated with identical secondary structure to several ITR arrays in circular intermediates. The primary sequence is completely altered and bears no resemblance to sequences contained within native AAV ITRs. These synthetic DNA sequences are tested for their ability to confer increased episomal stability to heterologous plasmids in several model systems including: 1) the airway, 2) muscle, 3) and developing *Xenopus* embryos. The developing *Xenopus* embryo model is ideal for testing integration and persistence of plasmid based vectors for application of *in utero* gene therapy. If synthetic DNA sequences with similar secondary structure to ITRs are found to confer increased persistence to plasmid based vectors, then determinants for protein binding which facilitate persistence are independent of primary base sequence. These studies allow the optimization of the secondary structural requirements by synthesizing a wide range of DNA molecules with varying degrees of palindromic repeats. Furthermore, the secondary structure may not bind proteins directly but facilitate recombination of plasmids to large concatamers which have increased episomal stability or enhanced integration efficiencies.

Example 5

Delivery of Multiple Genes through Intermolecular Concatamerization

Methods

Recombinant AAV vectors.

5 Two rAAV vector stocks were generated for use in these studies, AV.GFP3ori (Example 1) and AV.Alkphos (also known as CWRAPSP, a gift of Dusty Miller) (Halbert et al., 1997). Virus stocks were generated by co-transfection of 293 cells with either pCisAV.GFP3ori or pCWRAPSP along with pRep/Cap, followed by co-infection with recombinant Ad.CMVlacZ helper virus
10 (Example 2). rAAV was then purified through three rounds of CsCl density gradient centrifugation as previously described by Duan et al. (1997). Purified viral fractions were heated at 60°C for 1 hour to inactivate any residual contaminating helper adenovirus. The yields for AV.GFP3ori and AV.Alkphos were 1×10^{12} and 7×10^{11} particles per ml, respectively, as determined by slot
15 blot hybridization with ^{32}P -labeled GFP or Alkphos probes. Infectious titers determined by infection of 293 cells with rAAVs were 1.1×10^9 IU/ml (AV.GFP3ori) or 8.6×10^8 IU/ml (AV.Alkphos). Controls testing for contamination of rAAV stocks with wtAAV by anti-Rep immunocytochemical staining in rAAV/Ad.CMVlacZ co-infected 293 cells were negative (limit of
20 sensitivity is less than 1 infectious wtAAV particle per 10^{10} DNA particles of rAAV). Similarly, histochemical staining for β -galactosidase in rAAV infected 293 cells showed no detectable contamination with helper adenovirus in 10^{10} DNA particles of rAAV (limit of sensitivity).

Infection of muscle tissue and evaluation of transgene expression.

25 The C57BL/6 mice used for these experiments were housed in a virus-free animal care facility and were maintained under strict University of Iowa and NIH guidelines, using a protocol approved by the Animal Care and Use Committee and facility veterinarians. Four to five week old mice received bilateral 30 μl injections of a mixture of both AV.GFP3ori and AV.Alkphos into
30 the tibialis anterior muscle (5×10^9 DNA particles of each virus per muscle). Controls included uninjected muscles and muscles receiving injections of one of the viruses alone. At 14, 35, 80, and 120 days post-infection, animals were euthanized and tissues were harvested for evaluation of transgene expression and

preparation of low molecular weight Hirt DNA. For each experimental time point, at least 3 independently injected muscles were evaluated.

In all experiments, GFP fluorescence was visualized in freshly excised muscle tissue prior to processing. A portion of the same muscle was fixed with 2% paraformaldehyde in phosphate buffered saline, and cryoprotected in graded sucrose solutions before embedding in optimal cutting temperature medium (OCT). Sections (6 μ m) were then evaluated for GFP expression directly and Alkphos expression following heat inactivation of endogenous Alkphos and histochemical staining for Alkphos activity (Engelhardt et al., 1995). To confirm dual localization of GFP and Alkphos expression in the same muscle fibers, either serial sections were evaluated for GFP and Alkphos expression or the same section was first photographed for GFP expression followed by histochemical staining for Alkphos and re-imaging of the same field.

Rescue of circular intermediates from muscle Hirt DNA.

Low molecular weight Hirt DNA was prepared from 20 mg specimens of injected muscles from 3 animals at each time point (Example 2). Hirt DNA (4 μ l; 1/5 of the total volume) was then used to transform 50 μ l of electrocompetent SURE cells (Stratagene) using a BioRad *E. coli* electroporator and 0.1 μ m cuvettes. Colonies resulting from each bacterial transformation were quantified, and plasmids from 20 colonies from each muscle Hirt DNA sample were purified for analysis. It should be noted that only circular forms carrying the Amp resistance gene and the bacterial origin of replication from AV.GFP3ori are rescued by bacterial transformation (Duan et al., 1998). Control experiments reconstituting 5×10^{10} viral DNA particles into uninfected muscle extracts prior to Hirt DNA preparation failed to give rise to replication competent plasmids in the rescue assay (Duan et al., 1998). Additional controls in Duan et al. (1998) using AV.GFP3ori virus also demonstrated that linear double stranded and single stranded purified viral DNA genomes do not give rise to replication competent plasmids following transformation into *E. coli*.

Characterization of encoded genes in rescued circular intermediates.

Several assays were used to characterize the extent of intermolecular recombination between independent circular viral genomes by evaluating the number and type of encoded genes in rescued plasmids from Hirt DNA of

muscles co-infected with AV.GFP3ori and AV.Alkphos. Initial analysis involved the bulk evaluation of 60 rescued plasmids (20 from each of three muscle samples for each time point) by dot blot hybridization of mini-prep DNA with EGFP, Alkphos, and Amp ³²P-labeled DNA probes. In these studies, Amp hybridization served as a control to show that there was a sufficient quantity of DNA for the analysis. The percentages of Alkphos and/or GFP hybridizing plasmids were calculated by this method for each muscle sample. From this percentage, the total number of plasmids hybridizing to each probe in the Hirt DNA sample was calculated from the total CFU obtained in each transformation.

In this analysis, each muscle sample was evaluated independently to determine the mean (+/-SEM) total Alkphos and/or GFP hybridizing plasmids. A second evaluation involved the transfection of rescued plasmids into 293 cells using lipofectamine, followed by evaluation of GFP fluorescence and histochemical staining for Alkphos. To confirm that GFP and Alkphos co-expressing plasmids were indeed clonal and that both genes were encoded on the same plasmid, a selected group of five co-expressing plasmids were retransformed into *E.coli* and colonies were re-isolated prior to repeating the transfection studies. In all cases, plasmids co-expressing the two reporter genes remained clonal through this subsequent re-isolation.

Structural analysis of concatamer rAAV circular intermediates.

To further characterize the nature of isolated circular intermediates co-expressing both GFP and Alkphos transgenes, plasmid structure was mapped by Southern blotting and restriction enzyme analysis. The structural of five co-expressing circular intermediate plasmids were determined by digestion with AhdI, HindIII, NotI, HindIII/NotI, ClaI/AseI, and/or SnaBI and Southern blotting was performed with ³²P-labeled GFP, Alkphos, and ITR probes.

Results

Strategy for characterizing mechanisms of rAAV circular intermediate formation.

Efficient circularization of rAAV genomes has been previously demonstrated to occur in muscle in a time dependent fashion (Example 2). Furthermore, the conversion of monomeric to multimeric circular rAAV intermediates occurred over time and was associated with long-term episomal

persistence of AAV genomes. High molecular weight AAV circular genomes might form by either of the following two mechanisms, one involving the replication of monomer structures and the other through intermolecular recombination between independent monomers. A rescue assay was developed using two separate rAAV vectors, AV.GFP3ori and AV.Alkphos (Figure 14A), which allowed for the identification of independent viral genomes through unique transgenes. In this assay, circular form genomes were rescued in bacteria by virtue of Amp/ori sequences encoded in one of the two vectors (AV.GFP3ori). A method for characterizing the extent of intermolecular recombination between independent circular rAAV genomes was shown in Figure 14B.

Co-expression of independently encoded rAAV transgenes in muscle myofibers.

To confirm that myofibers can be co-infected at a high efficiency with the two rAAV vectors, the tibialis anterior muscle of mice was co-infected with 5×10^9 DNA particles of both AV.GFP3ori and AV.Alkphos. At 14, 35, 80, and 120 days post-infection, muscles were harvested and analyzed for transgene expression. Transgene expression from both reporters was weak but clearly visible in 14 day muscle samples. By 80 days post-infection, transgene expression was maximal and serial sections demonstrated expression of both Alkphos and GFP transgenes in overlapping regions of the muscle (Figures 15A-C). At this time point, approximately 50% of the fibers in the tibialis muscle expressed both transgenes. To confirm that co-infection of myofibers occurred with the two independent vectors, co-localization studies were performed on muscle sections by a serial staining procedure. These studies, depicted in Figure 15D, demonstrate four classes of myofiber transgene expression: 1) GFP positive only, 2) Alkphos positive only, 3) GFP/Alkphos positive, and 4) no transgene expression. The largest fraction of myofibers expressed both GFP and Alkphos transgenes. These results confirm that at the titers of virus used for infection, co-infection occurred in greater than 90% of transgene expressing myofibers.

Rescue of bi-functional rAAV circular intermediates increases over time.

To determine the extent of recombination between circular AAV genomes, circular form genomes were rescued as plasmids from low molecular weight Hirt DNA of muscle tissue co-infected with AV.GFP3ori and

AV.Alkphos. Following transformation of *E.coli* Sure cells with Hirt DNA purified from infected muscles, the total number of GFP and Alkphos hybridizing Amp resistant bacterial plasmids was quantitated for each time point post-infection (Figure 16A and B) (Duan et al., 1995), the abundance of circular AAV genomes rescued from AV.GFP3ori increased over time. For each muscle sample (three for each time point) twenty plasmid clones were evaluated for hybridization to GFP and Alkphos DNA probes and the total number of plasmids was back calculated from the total CFU for each individual muscle sample. Figure 16B demonstrates the mean (\pm SEM, N=3) total plasmids that hybridized to GFP or GFP/Alkphos probes at each time point. At 14 days post-infection, GFP/Alkphos co-hybridizing plasmids were never observed. In contrast, at time points after 35 days the percentage of GFP/Alkphos co-hybridizing plasmids increased with time and reached 33% by 120 days (Figure 16C). Since bacterial plasmid rescue can only occur through AV.GFP3ori genomes, this data suggests that recombination between independent Alkphos and GFP rAAV genomes takes place over time. These results are consistent with studies described hereinabove demonstrating a time dependent concatamerization of monomer circular rAAV genomes in muscle.

To evaluate the ability of circular intermediates to express encoded transgenes, transient transfection studies were performed in 293 cells with rescued circular intermediate plasmids (Figures 17A-C). Between 85-90% of rescued plasmids hybridizing to GFP probes on slot blots also expressed the GFP transgene in this transfection assay (Figure 17D). The percentage of GFP expressing plasmids that also expressed Alkphos rose over time in concordance with the hybridization data (Figure 17D). However, approximately 40-50% of plasmids which were hybridization positive for Alkphos did not express the Alkphos transgene. This may represent recombinational deletion of the RSV promoter driving Alkphos expression which occurred during concatamerization at sites near the 5' ITR. These results demonstrate that intermolecular recombination between Alkphos and GFP derived circular intermediates occurs as part of the time dependent concatamerization process of rAAV in muscle. To confirm that amplified plasmids stocks expressing both reporter genes were actually clonal (i.e., one plasmid rather than two independent plasmids resulting

from contamination), a select number of bacterial clones expressing both transgenes were re-isolated and the transfection assays were repeated. In all cases, plasmids expressing the two reporter genes remained clonal through two rounds of bacterial cloning. Hence, dual reporter expression was not due to
5 contamination of independent GFP and Alkphos expressing plasmids.

Concatamerization of AAV circular intermediates occurs through uniform intermolecular recombination between ITRs of independent viral genomes.

To better understand the mechanisms of circular concatamer formation, a detailed structural analysis was performed of five bi-functional circular
10 concatamers isolated from rAAV infected muscle samples. As previously described for the AV.GFP3ori genome (Example 2), the conversion of monomeric circular AAV genomes to large multimeric circular concatamers with a predominant head-to-tail structure increased with time in muscle. To evaluate the structure of bi-functional circular concatamers, restriction enzyme mapping
15 and Southern blot analysis using ³²P-labeled EGFP, Alkphos, and ITR probes was employed. Results from five analyzed plasmids demonstrated between 3-6 genomes within these circular concatamers. Two representative structures from 35 and 80 day time points are shown in Figure 18. Several interesting conclusions can be made from this structural analysis. As described, head-to-
20 tail oriented genomes could be seen in all isolated concatamers. However, several examples of head-to-head and tail-to-tail genome combinations of AV.Alkphos and AV.GFP3ori were also seen. Since head-to-head and tail-to-tail genome concatamers were never seen in muscles infected with AV.GFP3ori alone, there must be a selective disadvantage for bacterial replication when ori
25 sequences are in either of these conformations. However, since the AV.Alkphos genomes do not contain a bacterial origin of replication, this orientation is permitted in chimeric concatamers. Second, noticeable deletions and/or loss of restriction sites close to ITRs were noted (Figure 17). It is not known whether deletions close to the ITR are a common event in the concatamerization process,
30 but if so, this could account for the fact that only 60% of GFP/Alkphos hybridizing circular intermediates also expressed the Alkphos transgene.

Discussion

Concatamerization of rAAV genomes has long been recognized in integrated proviral genomes. Recently, the association of this concatamerization process with the formation of high molecular circular genomes in muscle has suggested that this process may also be important in episomal persistence. The findings described herein demonstrated rescue of independent viral genomes within the same circular concatamer, suggesting that this process of concatamerization occurs through intermolecular recombination. Furthermore, at 14 days the predominant form of viral genome in muscle was circular monomers (Example 2), which correlates with the results described above demonstrating only GFP expression in rescued circular intermediates at this time point. Together with the fact that bi-functional rescued circular concatamers increase with time, these results suggest that large concatamers form by recombination of monomeric circular precursor genomes. Furthermore, since an alternative model of concatamerization by rolling circular replication would be expected to yield only GFP expressing rescued plasmids in this system, this mechanism does not appear responsible for concatamerization.

Based on the structural analysis of these bi-functional circular intermediates, recombination between monomeric circular rAAV genomes is likely facilitated through ITR sequences. Directionality of this recombinational event does not appear to play a significant role, since head-to-tail, head-to-head, and tail-to-tail oriented intermolecular concatamers were found. In addition, the extent to which recombination within ITR repeat regions occurs in bacteria is presently unknown and may account for the deletions and/or restriction site losses near ITR arrays. However, serial passaging of bi-functional circular AAV genomes in bacteria has suggested that the structure of these large concatamers is impressively stable in bacteria.

Intermolecular recombination of rAAV genomes to form single circular episomes may be particularly useful for gene therapy. For example, large regulatory elements and genes beyond the packaging capacity of rAAV may become linked after co-infecting tissue with two independent vectors (Figure 19). This strategy could also involve trans-splicing vectors encoding two independent regions of a gene which are brought together to form an intact

splicing unit by circular concatamerization.

Example 6

Enhancement of Recombinant AAV Mediated Gene Expression

Through Intermolecular *Cis*-activation

Materials and Methods

Recombinant AAV vectors.

The pcisAV.Luc proviral plasmid was generated by cloning the 1983 bp NheI/BamHI fragment from pGL3-Basic (Promega), containing the luciferase gene and SV40 late polyA signal, by blunt-end ligation into the blunted Xba I site of pSub201 (Samulski et al., 1987). Similarly, pcisAV.SV(P)Luc was generated using a blunted 2175 bp NheI/BamHI fragment, from the pGL3-Promoter (Promega), containing the SV40 promoter, luciferase gene, and SV40 late polyA signal. The pcisAV.SV(P/E)Luc plasmid was generated by blunt-end ligation of a 2427 bp NheI/SalI fragment from pGL3-Control (Promega) into the blunted Xba I site of psub201. This construct contains the SV40 promoter, luciferase gene, SV40 late polyA signal and SV40 enhancer.

The "super-enhancer" vector, pcisAV.SupEnh, was produced using a two-step cloning process. First, a 0.62 kb blunted BglII/PvuI fragment containing the CMV immediate early enhancer from pIRES (Clontech) was subcloned into the blunted BamHI site in pGL3-Control (Promega) to make pGL3-Control-CMVenh. Then a 0.92 kb DNA segment containing both the CMV immediate early enhancer and the SV40 enhancer was released by ClaI/SalI double digestion of pGL3-Control-CMVenh and subsequently inserted into the blunted PstI site of pcisAV.GFP3ori (Duan et al., 1998b). The resulting pcisAV.SupEnh plasmid contains the SV40 enhancer, the CMV immediate early enhancer, the β -lactamase gene, and a bacterial replication origin. The ampicillin resistance gene (β -lactamase) and bacterial original of replication were included in pcisAV.SupEnh to facilitate the subsequent rescue of circular AAV genomes from infected cells in bacteria.

The control vector, pcisAV.AmpOri, was generated by blunt-end ligation of a 1.1 kb SalI digested stuffer sequence from the human glycosylasparaginase cDNA into PstI digested pcisAV.GFP3ori. This plasmid has a structure similar

to that of pcisAV.SupEnh, except that it does not contain any enhancer elements. The pcisAV.AmpOri was used as a negative control for non-specific enhancement of transgene expression by intermolecular recombination of two different AAV vectors.

5 The integrity of the ITR sequences in all the plasmids was confirmed by digestion with restriction enzymes, including SmaI, MscI, and BssHII, which have unique cutting sites within different regions of ITR. All the viral preparations were obtained according to a method described in Duan et al. (1997). The quality of the viral stocks (i.e., contamination with adenovirus
10 and/or wild type AAV) was confirmed as previously described in Duan et al. (1998b). The analyses showed less than 1 recombinant adenovirus and wt AAV infectious particles per 10^{10} particles of rAAV. Viral titers were determined by quantitative slot-blot hybridization using either luciferase, CMV enhancer, SV40 enhancer, or *ori* probes for each of the respective vectors against plasmid copy
15 number standards.

Luciferase enzyme assays.

Luciferase assays were performed from cell lysates harvested from either *in vitro*-infected human fibroblasts or from *in vivo*-infected mouse tibialis anterior muscle. Human fibroblasts were infected with virus in 60 mm dishes.
20 These *in vitro*-infected cells were harvested at 3 days post-infection by rinsing cells with PBS twice, and then incubating with 1x Report lysis buffer (Promega) (400 μ l per 60 mm plate) at room temperature for 15 minutes. Cells were scraped into an eppendorf tube and centrifuged for 30 seconds at 14,000 rpm. Serial dilutions of supernatants were assayed for luciferase activity using
25 reagents from Promega according to the manufacturer's protocols. Luciferase activity was detected in triplicate for each individual sample with a Luminometer (TD-20/20, Turner Designs Instrument, Sunnyvale, CA), at the setting of 90.3% sensitivity, 1 second delay, 10 second measurement. Six independent samples were assayed for each experimental condition.

30 For *in vivo* assay of luciferase activity, the anterior tibialis muscle of 8 week old C57BL/6 mice was infected with an indicated amount and type of AAV vectors (see Figure 23) in 30 μ l phosphate-buffered saline (PBS). The entire muscle was harvested at 30 days or 90 days post-infection and weighed

prior to cell lysate preparation. The muscle tissue was frozen in liquid nitrogen and pulverized by hand grinding with an ice-cold porcelain mortar and pestle. The muscle was further minced and homogenized in 100 µl of 1x Report lysis buffer with a hand-held plastic pestle for 2 minutes (Kontes, Vineland, New Jersey). After 15 minutes incubation at room temperature, the crude lysates were spun for 30 seconds at 14,000 rpm, and the supernatants were used for luciferase activity assay as described above. To minimize variability, all experimental samples were analyzed simultaneously using the same batch of luciferase assay reagents and were normalized to the protein content in the lysate.

10 Results

Co-administration of a *cis*-activating vector increases rAAV mediated luciferase expression in fibroblasts.

To test the hypothesis that *cis*-activation from two independent AAV vectors can occur following intermolecular concatamerization, several rAAV vectors with defined regulator elements and/or the luciferase reporter gene were constructed (Figure 20). One of these vectors, AV.Luc, contains the luciferase transgene and an SV40 poly A signal but no promoter sequences. AV.SV(P)Luc is similar to AV.Luc except that an SV40 promoter (lacking the enhancer sequence) was inserted in front of the luciferase transgene. AV.SV(P/E)Luc includes both the SV40 promoter and enhancer, driving expression of the luciferase transgene, and was used as a control for maximal expression in the absence of intermolecular recombination with an enhancer containing vector. To evaluate intermolecular *cis*-activation by enhancer elements, an rAAV “super-enhancer” vector (AV.SupEnh) was also constructed, which contains SV40 and CMV enhancer regions without promoter or transgene sequences. A negative control vector (AV.AmpOri), which is similar to AV.SupEnh except that the enhancer sequences were replaced by a non-specific stuffer fragment, was also constructed.

Enhancers are *cis*-acting DNA sequences that can be recognized by regulatory proteins to stimulate transcription in a context independent manner relative to the promoter and transgene. If cells were co-infected with AV.SV(P)Luc and AV.SupEnh vectors, luciferase transgene expression could be significantly increased from the minimal SV40 promoter only if intermolecular

recombination had occurred between the two independent vectors (Figure 21). However, in accordance with the definition of an enhancer, no activation should occur if the enhancer sequences and the transgene cassette are located in separate circular DNA molecules (Lewin, 1997).

5 Initial experiments were performed by infecting 1×10^6 primary human fibroblasts with single vectors [AV.Luc or AV.SV(P)Luc] at a multiplicity of infection (moi) equal to 1000 viral particles/cell. Additional experimental points included co-infection of AV.Luc or AV.SV(P)Luc with either AV.SupEnh or AV.AmpOri at the same moi.

10 As shown in Figure 22, infection of fibroblasts with both the promoter-less [AV.Luc] and the minimal promoter [AV.SV(P)Luc] luciferase constructs alone gave only minimal expression at 3 days post-infection. However, co-infection with AV.SupEnh produced 16- and 35-fold inductions in luciferase expression from the AV.Luc and AV.SV(P)Luc vectors, respectively. Thus, *cis*-
15 activation of a minimal SV40 promoter can occur in the presence of a second AAV vector containing enhancer elements. Unanticipated, however, was the high level of transactivation of the AV.Luc construct, which contains no heterologous promoter sequences. These findings support earlier studies suggesting that ITRs contain a cryptic promoter (Flotte et al., 1993). The
20 specificity of this induction was further demonstrated by the lack of transactivation following co-administration of an alternative control vector, AV.AmpOri, which does not contain enhancer elements.

To confirm that the transactivation observed was due to recombination of two independent AAV virus, Hirt DNA from infected cells was transformed into
25 competent SURE *E. Coli* cells. As expected, no bacterial clones were retrieved from cells infected with either AV.Luc or AV.SV(P)Luc alone (neither vector contains *amp*^r and *ori* sequences). However, in cells co-infected with AV.SupEnh (which contains *amp*^r and *ori* sequences but no luciferase gene), approximately 4% of the rescued clones also contained the luciferase transgene,
30 according to restriction enzyme mapping and Southern blotting analyses. Subsequent transfection of Hela cells with these rescued, circular concatamer plasmids demonstrated 103 \pm 6 fold higher luciferase activity from AV.Luc/AV.SupEnh, as compared to AV.Luc/AV.AmpOri recombined AAV

genome plasmids. Taken together, these results indicated that intermolecular concatamerization of a reporter rAAV virus with an independent "super-enhancer" AAV vector substantially increased the efficiency of transgene expression in cultured human primary fibroblasts.

5 Intermolecular *cis*-activation enhances AAV mediated luciferase expression in muscle tissue *in vivo*.

To confirm whether the *in vitro* findings could also be applied to increase rAAV mediated gene expression *in vivo*, 2×10^{10} particles of AV.Luc or AV.SV(P)Luc were injected into the tibialis anterior muscle of C57BL/6 mice,
10 either individually or in combination with AV.SupEnh (2×10^{10} particles). The infected muscles were harvested at 30 and 90 days post-infection. Consistent with previous findings (Example 5), intermolecular recombination of circular rAAV genomes increased from about 5% at the 30 day time point to 25% by 90 days post-infection. Importantly, co-administration of AV.SupEnh vector with
15 either AV.Luc or AV.SV(P)Luc (Figure 23A) led to a functional enhancement of transgene gene expression, with a 30-fold increase observed at 30 days post-infection. By 90 days post-infection, greater than 200-fold [AV.Luc] and 600-fold [AV.SV(P)Luc] increases in transgene expression were observed when AV.SupEnh was co-administered.

20 Consistent with *in vitro* experiments in fibroblasts, the bacterially rescued concatamers containing both the luciferase transgene and AV.SupEnh also demonstrated greater than a 100-fold higher luciferase activity than the original proviral luciferase plasmid (pcisAV.Luc) alone. Interestingly, 90 day muscles infected with 2×10^{10} particles of the control AV.SV(P/E)Luc vector
25 (which contains both the SV40 enhancer and promoter) produced luciferase levels that were 10- and 100-fold less than the levels seen following co-infection with AV.SupEnh/AV.Luc and AV.SupEnh/AV.SV(P)Luc, respectively. These findings underscore the potential of using large multi-enhancer segments to increase AAV mediated gene delivery through intermolecular *cis*-activation.
30 Such applications will likely have broad implications for *in vivo* AAV mediated gene delivery to organs for which concatamerization of the AAV genome is an inherent process of its latent life cycle.

Both *in vitro* and *in vivo* results described herein demonstrated effective augmentation of transgene expression following intermolecular concatamerization between two independent AAV vectors carrying a transgene or enhancer sequences. A surprising finding was the strong induction of transgene expression from an AAV vector lacking an endogenous promoter (AV.Luc). Previous studies have implied weak promoter activity in the AAV ITR sequences (Flotte et al., 1993). Hence, it is possible that, in combination with enhancer sequences supplied by another AAV vector, a therapeutic level of transgene expression could be achieved for disease genes approaching the maximum packaging capacity for AAV. One notable example would be in applications of AAV for cystic fibrosis. Additionally, a particular interesting extension of this work will be to use large cell-specific enhancer regions for targeted expression.

15

Example 7

Generation of rAAV Delivery Systems for Full-length CFTR and EPO

Based on AAV Circular Concatamer Formation

Given the findings that circularization and concatamerization are integral parts of rAAV transduction in the airway, rAAV transduction, circularization and concatamerization can be used to deliver CFTR transgene cassettes to the airway. Two approaches may be employed in which various genetic elements of the CFTR transgene cassette are split into two or more vectors which are then used for co-infection of the airway. The first approach, "*trans*-splicing", utilizes a rAAV vector harboring the promoter/enhancer driving the first half of CFTR DNA flanked by a donor splice site (vector-1, Figure 19C) and a second rAAV vector harboring the second half of CFTR DNA and a polyA addition sequence preceded by a donor acceptor site (vector-2, Figure 19C). A second approach, "*cis*-activation", employs a first vector that harbors the entire CFTR transgene with a minimal synthetic promoter and a second vector comprises several strong enhancer sequences. Through the process of concatamerization, these two vectors are brought into juxtaposition with one another, allowing for splicing or

cis-activation of enhancer/promoter combinations.

Initial experiments assess the extent of concatamerization in epithelia and in cell lines. Two rAAV CFTR plasmid-based vectors were prepared, pAV.RVSCFTR donor and pAV.CFTR acceptor. The splice consensus sequences are based on the large T antigen splice sites. A synthetically generated circular chimera was also constructed to represent rAAV circular concatamers. The chimera was generated by PCR-mediated approaches in which splice site consensus sequences were incorporated into the primers used to amplify each half of CFTR from pBQCFTR (Drumm et al., 1990). Several clones were sequenced in their entirety and PCR errors removed by subcloning the appropriate correct fragment from pBQCFTR. Assays for splicing integrity include RNase protection assays of transfected MDCK cells for sequences within the large T antigen intron and surrounding regions. Polarized airway epithelial cells grown at the air-liquid interface are co-infected with the donor and acceptor CFTR AAV vectors and CFTR gene expression in these cells is then monitored by both immunofluorescent localization and functional analysis of short circuit currents (Smith et al., 1992; Smith et al., 1990). Additionally, functional assays using two electrode voltage clamp measurements (TEV) of oocytes following nuclear microinjection of the chimeric plasmid are used to demonstrate intact splicing and active CFTR channels (Zhang et al., 1998). Hirt analyses of episomal AAV species are used to correlate the efficacy and persistence of CFTR gene expression with the formation of AAV circular intermediates.

Several systems are utilized to evaluate the efficiency of intermolecular recombination to form functional CFTR expressing circular concatamers. The first employs short circuit measurements in polarized CF airway epithelia. Conditions for rAAV infection include basolateral infection of polarized CF epithelia, preferably in the presence of agents that enhance the level of transgene expression. A second model employs CF human bronchial xenografts to evaluate CFTR function complementation. Reconstitution of xenografts with UV-irradiated rAAV infected primary cells can result in approximately 50% transduction in differentiated xenografts. Functional expression of CFTR in CF primary airways is evaluated by transepithelial potential difference (PD)

measurements (Jiang et al., 1998):

In addition, a "cis-activation" approach may be employed (Example 6) to utilize the concatamerization process to deliver full-length CFTR. For example, a synthetic minimal promoter driving full-length CFTR within one viral vector and a second independent vector encoding a tandem array of strong enhancer sequences such as RSV, CMV, and SV40 may be employed.

A trans-splicing approach was employed to deliver the genomic epo gene to mice. The vectors are shown in Figure 25. The infection of mice with these vectors protected the mice from adenine induced anemia brought on by renal failure.

Example 8

Generation of rAAV Delivery Systems for Autonomously Replicating Episomes Harboring EBV Replication Origins

To prepare autonomously replicating circular episomes, e.g., to increase the stability of genes delivered via rAAV vectors, a rAAV vector comprising a replication origin of a circular episome is employed. For example, a rAAV vector comprising the EBV replication origin (OriP) and EBNA-1, the only viral protein needed to facilitate replication at this origin, is prepared (Vector 2, Figure 24). DNA fragments encoding EBNA-1 and OriP are excised from pREP10 (Clontech Inc.) and cloned into a pCisAV.RSV vector. Initially, virus is produced from pCisAV.EBNA-1/OriP construct and a reporter construct, e.g., AV.GFP3ori, and virus is used to co-infect Hela cells. Hela cells are infected with each of the vectors alone (Figure 24) or together (MOI = 1000 particles/cell) and serially passaged. Transgene expression is quantitated following each passage and Hirt DNA is isolated for Southern blot analysis of episomal DNA persistence. If persistence of GFP expression is increased in the presence of co-infection with EBNA-1 rAAV vectors following serial passage, there will be an increased abundance of co-encoding EBNA-1/GFP rescued plasmids with increasing passage number.

Then primary human bronchial cells are infected with each of the vectors alone or together at an MOI = 10,000 particles/cell and infected cells seeded into

human bronchial xenografts. Persistence of GFP transgene expression is compared between infected xenografts at 1, 3, 5, and 8 weeks post-transplantation. Epithelia is also harvested for Hirt DNA analysis by perfusion of xenografts with SDS containing Hirt extraction buffer. Hirt DNA Southern
5 blots using EBNA and GFP probes assess the extent of episomal persistence of monomer and concatamer circular genomes over time.

An increase in the persistence of EBNA-1 hybridizing Hirt DNA in cells infected with the EBNA-1 vector alone, as compared to GFP-probed Hirt Southern of cells infected with only the GFP encoding rAAV construct, is
10 observed as monomer circularized EBNA-1 vectors are the predominant form of autonomously replicating episomes. However, when cells are infected with both vectors simultaneously, GFP transgene expression is prolonged. Molecular analyses are performed to characterize the autonomously replicating circular chimeras encoding both GFP and EBNA-1.

15 This approach may be employed for gene therapy, as circular concatamers with as many as five genomes, have been observed. For example, for CFTR gene therapy, human bronchial xenografts and the RSV-EBNA-1 rAAV vector and a second CFTR rAAV vector with a minimal promoter may be employed. If genomes are oriented in a head-to-head fashion, the RSV enhancer
20 may increase transcription from the minimal CFTR promoter.

References

- Afione, S.A., Conrad, C.K., Kearns, W.G., Chunduru, S., Adams, R., Reynolds, T.C., Guggino, W.B., Cotting, G.R., Carter, B.J., and Flotte, T.R. *In vivo*
25 model of adeno-associated virus vector persistence and rescue. *J. Virol.* 70, 3235-3241 (1996).
- Ali, R.R. *et al.* Gene transfer into the mouse retina mediated by an adeno-associated viral vector. *Hum Mol Genet* 5, 591-594 (1996).
- Bennett, J., Duan, D., Engelhardt, J.F., and Maguire, A.M. Real-time
30 noninvasive *in vivo* assessment of adeno-associated virus-mediated retinal transduction. *Invest. Ophthalmol. Vis. Sci.* 38, 2857-2963 (1997).
- Berns, K.I. Parvovirus replication. *Microbiol Rev* 54, 316-329 (1990).
- Berns, K.I. & Giraud, C. Biology of adeno-associated virus. *Curr. Top.*

Microbiol Immunol. 218, 1-23 (1996).

Cheung, A.K., Hoggan, M.D., Hauswirth, W.W. & Berns, K.I. Integration of the adeno-associated virus genome into cellular DNA in latently infected human Detroit 6 cells. *J. Virol.* 33, 739-748 (1980).

- 5 Clark, K.R., Sferra, T.J. & Johnson, P.R. Recombinant adeno-associated viral vectors mediate long-term transgene expression in muscle. *Hum. Gene Ther.* 8, 659-669 (1997).

- Conrad, C.K., Allen, S.S., Afione, S.A., Reynolds, T.C., Beck, S.E., Fee-Maki, M., Barraza-Ortiz, X., Adams, R., Askin, F.B., Carter, B.J., Guggino, W.B., and Flotte, T.R. Safety of single-dose administration of an adeno-associated virus (AAV)-CFTR vector in the primate lung. *Gene Ther.* 3, 658-668 (1996).
- 10

- Drumm, M.L., Pope, H.A., Cliff, W.H., Rommens, J.M., Marvin, S.A., Tsui, L.C., Collins, F.S., Fritzzell, R.A. & Wilson, J.M. Correction of the cystic fibrosis defect in vitro by retrovirus-mediated gene transfer. *Cell* 62, 1227-1233 (1993).
- 15

Duan, D., Fisher, K.J., Burda, J.F. & Engelhardt, J.F. Structural and functional heterogeneity of integrated recombinant AAV genomes. *Virus Res* 48, 41-56 (1997).

- 20 Duan, D. *et al.* Circular Intermediates of Recombinant Adeno-Associated Virus have Defined Structural Characteristics Responsible for Long Term Episomal Persistence In Muscle. *J. Virol.* 72, 8568-8577 (1998b).

- Duan, D., Yue, Y., Yan, Z., McCray, P.B. & Engelhardt, J.F. Polarity Influences the efficiency of recombinant adeno-associated virus infection in differentiated airway epithelia. *Human Gene Therapy* 9, 2761-2776 (1998a).
- 25

Duan, D., Yan, Z., Yue, Y. & Engelhardt, J.F. Structural analysis of adeno-associated virus transduction intermediates. *Virology* 261, 8-14 (1999).

- Engelhardt, J.F., Schlossberg, H., Yankaskas, J.R. & Dudus, L. Progenitor cells of the adult human airway involved in submucosal gland development. *Development* 121, 2031-2046 (1995).
- 30

Ferrari, F.K., Samulski, T., Shenk, T., and Samulski, R.J. Second-strand synthesis is a rate-limiting step for efficient transduction by recombinant

- adeno-associated virus vectors. *J. Virol* 70, 3227-3234 (1996).
- Fisher, K.J., Gao, G.P., Weitzman, M.D., DeMatteo, R., Burda, J.F., and Wilson, J.M. Transduction with recombinant adeno-associated virus for gene therapy is limited by leading-strand synthesis. *J. Virol.* 70, 520-532 (1996).
- 5 Fisher, K.J. et al. Recombinant adeno-associated virus for muscle directed gene therapy. *Nat. Med.* 3, 306-312 (1997).
- Fisher-Adams, G., Wong, K.K., Jr., Podsakoff, G., Forman, S.J. & Chatterjee, S. Integration of adeno-associated virus vectors in cd34+ human hematopoietic progenitor cells after transduction. *Blood* 88, 492-504 (1996).
- 10 Flannery, J.G. et al. Efficient photoreceptor-targeted gene expression *in vivo* by recombinant adeno-associated virus. *Proc. Natl. Acad. Sci. USA* 84, 6916-6921 (1997).
- 15 Flotte, T.R. et al. Expression of the cystic fibrosis transmembrane conductance regulator from a novel adeno-associated virus promoter. *J Biol Chem* 268, 3781-3790 (1993).
- Flotte, T.R., Afione, S.A. & Zeitlin, P.L. Adeno-associated virus vector gene expression occurs in nondividing cells in the absence of vector DNA integration. *Amer. J. Resp. Cell Mol. Biol.* 11, 517-521 (1994).
- 20 Halbert, C.L. et al. Transduction by adeno-associated virus vectors in the rabbit airway: Efficiency, persistence and readministration. *J. Virol.* 71, 5932-5941 (1997).
- Herzog, R.W., Hagstrom, J.N., Kung, S.H., Tai, S.J., Wilson, J.M., Fisher, K.J., and High, K.A. Stable gene transfer and expression of human blood coagulation factor IX after intramuscular injection of recombinant adeno-associated virus. *Proc. Natl. Acad. Sci. USA* 94, 5804-5809 (1997).
- 25 Hirt, B. Selective extraction of polyoma DNA from infected mouse cell culture. *J. Mol. Biol.* 26, 365-369 (1967).
- 30 Hyde, H., Davies, A.A., Benso, F.E., West, S.C. Resolution of recombination intermediates by a mammalian activity functionally analogous to *Escherichia coli* RuvC resolvase. *J. Biol. Chem.* 269, 5202-5209 (1994).

- Jiang, Q., Mak, D., Devidas, S., Schweibert, E.M., Bragin, A., Zhang, Y., Skach, W.R., Guggino, W.B., Foskett, J.K., and Engelhardt, J.F. Cystic fibrosis transmembrane conductance regulator-associated ATP release is controlled by a chloride sensor. *J. Cell Biol.* 143(3):645-57.
- 5 Kaplitt, M.G., Leone, P., Samulski, R.J., Xiao, X., Pfaff, D.W., O'Malley, K.L., and During, M.J. Long-term gene expression and phenotypic correction using adeno-associated virus vectors in the mammalian brain. *Nat. Genet.* 8, 148-154 (1994).
- Kearns, W.G. et al. Recombinant adeno-associated virus (AAV-CFTR) vectors
10 do not integrate in a site-specific fashion in an immortalized epithelial cell line. *Gene Ther.* 3, 748-755 (1996).
- Kessler, P.D. et al. Gene delivery to skeletal muscle results in sustained expression and systemic delivery of a therapeutic protein. *Proc. Natl. Acad. Sci. USA* 93, 14082-14087 (1996).
- 15 Koeberl, D.D., Alexander, I.A., Halbert, C.L., Russell, D.W. & Miller, A.D. Persistent expression of human clotting factor IX from mouse liver after intravenous injection of adeno-associated virus vectors. *Proc. Natl. Acad. Sci. USA* 94, 1426-1431 (1997).
- Kotin, R.M., Linden, R.M. & Berns, K.I. Characterization of a preferred site on
20 human chromosome 19q for integration of adeno-associated virus DNA by non-homologous recombination. *Embo J.* 11, 5071-5078 (1992).
- Lee, J., Voziyanov, Y., Pathania, S., Jayaram, M. Resolution of recombination intermediates by a mammalian activity functionally analogous to *Escherichia coli* RuvC resolvase. Structural alterations and
25 conformational dynamics in Holliday junctions induced by binding of a site-specific recombinase. *Mol. Cell* 1, 483-493 (1998).
- Lewin, B. *Genes VI*, xviii, 1260 (Oxford University Press, Oxford England ; New York, 1997).
- Linden, R.M., Ward, P., Giraud, C., Winocour, E., and Berns, K.I. Resolution of
30 recombination intermediates by a mammalian activity functionally analogous to *Escherichia coli* RuvC resolvase. Site-specific integration by adeno-associated virus. *Proc. Natl. Acad. Sci. USA* 93, 11288-11294 (1996a).

- Linden, R.M., Winocour, E. & Berns, K.I. The recombination signals for adeno-associated virus site specific integration. *Proc Natl. Acad. Sci. USA* 93, 7966-7972 (1996b).
- 5 Lobel, L.I., Murphy, J.E., and Goff, S.P. Resolution of recombination intermediates by a mammalian activity functionally analogous to *Escherichia coli* RuvC resolvase. The palindromic LTR-LTR junction of Moloney murine leukemia virus is not an efficient substrate for proviral integration. *J. Virol.* 63, 2629-2637 (1989).
- 10 McLaughlin, S.K., Collis, P., Hermonat, P.L. & Muzyczka, N. Adeno-associated virus general transduction vectors: Analysis of proviral structures. *J. Virol.* 62, 1963-1973 (1988).
- Miao, C.H. *et al.* The kinetics of rAAV integration in the liver [letter]. *Nat Genet* 19, 13-15 (1998).
- 15 Muzyczka, N. Use of adeno-associated virus as a general transduction vector for mammalian cells. *Curr. Top. Microbiol. Immunol.* 158, 97-129 (1992).
- Panganiban, A.T. Retroviral DNA integration. *Cell* 42, 5-6 (1985).
- Phillip, R. *et al.* Efficient and sustained gene expression in primary T
- 20 lymphocytes and primary and cultured tumor cells mediated by adeno-associated virus plasmid DNA complexed to cationic liposomes. *Mol. Cell. Biol.* 14, 2411-2418 (1994).
- Ponnazhagan, S., Erikson, D., Kearns, W.G., Zhou, S.Z., Nahreini, P., Wang, X.S., and Srivastava, A. Lack of site-specific integration of the
- 25 recombinant adeno-associated virus 2 genomes in human cells. *Hum. Gene Ther.* 8, 275-284 (1997).
- Qing, K., Khuntirat B., Mah C., Kube D.M., Wang X., Ponnazhagan, S., Zhou S., Dwarki V.J., Yoder M.C., Srivastava, A. Adeno-associated virus type 2-mediated gene transfer: correlation of tyrosine phosphorylation of the
- 30 cellular single-stranded D sequence-binding protein with transgene expression in human cells in vitro and murine tissues in vivo. *J. Virol.* 72, 1593-1599 (1998).
- Qing, K. *et al.* Role of tyrosine phosphorylation of a cellular protein in adeno-

- associated virus 2-mediated transgene expression. *Proc Natl Acad Sci U S A* 94, 10879-10884 (1997).
- Qing, K. *et al.* Adeno-associated virus type 2-mediated gene transfer: correlation of tyrosine phosphorylation of the cellular single-stranded D sequence-binding protein with transgene expression in human cells in vitro and murine tissues in vivo. *J Virol* 72, 1593-1599 (1998).
- Qing, K. *et al.* Human fibroblast growth factor receptor 1 is a co-receptor for infection by adeno-associated virus 2. *Nat Med* 5, 71-77 (1999).
- Reich, N.C., Sarnow, P., Duprey, E., and Levine, A.J. (1983). Monoclonal antibodies which recognize native and denatured forms of the adenovirus DNA-binding protein. *Virology* 128, 480-484.
- Rutledge, E.A. & Russell, D.W. Adeno-associated virus vector integration junctions. *J. Virol.* 71, 8429-8436 (1997).
- Samulski, R.J., Chang, L.S. & Shenk, T. A recombinant plasmid from which an infectious adeno-associated virus genome can be excised *in vitro* and its use to study viral replication. *J. Virol.* 61, 3096-3101 (1987).
- Samulski, R.J., Chang, L.S. & Shenk, T. Helper-free stocks of recombinant adeno-associated viruses: normal integration does not require viral gene expression. *J. Virol.* 63, 3822-3828 (1989).
- Samulski, R.J. Adeno-associated virus: integration at a specific chromosomal locus. *Curr. Opin. Genet. Dev.* 3, 74-80 (1993).
- Smith J.J., et al. cAMP stimulates bicarbonate secretion across normal, but not cystic fibrosis airway epithelia. *J. Clin Invest.* 89(4):1148-53 (1992).
- Smith J.J., et al. Bradykinin stimulates airway epithelial Cl-secretion via two second messenger pathways. *Am. J. Physiol.* 258:L369-77 (1990).
- Summerford, C. & Samulski, R.J. Membrane-associated heparan sulfate proteoglycan is a receptor for adeno-associated virus type 2 virions. *J. Virol.* 72, 1438-1445 (1998).
- Summerford, C., Bartlett, J.S. & Samulski, R.J. AlphaVbeta5 integrin: a co-receptor for adeno-associated virus type 2 infection. *Nat Med* 5, 78-82 (1999).

- Synder, R.O., MIAO, C.H., Patijn, G.A., Spratt, S.K., Danos, O., Nagy, D., Gown, A.M., Winther, B., Meuse, L., Cohen, L.K., Thompson, A.R., Kay, M.A. Persistent and therapeutic concentration of human factor IX in mice after hepatic gene transfer of recombinant AAV vectors. *Nat. Genet.* 16, 270-276 (1997).
- Varmus, H.E. Form and function of retroviral proviruses. *Science* 216, 812-820 (1982).
- Vieweg, J. et al., Efficient gene transfer with adeno-associated virus-based plasmids complexed to cationic liposomes for gene therapy of human prostate cancer. *Cancer Res.* 55, 2366-2372 (1995).
- Vincent-Lacaze, N. et al. Structure of adeno-associated virus vector DNA following transduction of the skeletal muscle *J Virol* 73, 1949-1955 (1999).
- Walsh, C.E., Nienhuis, A.W., Samulski, R.J., Brown, M.G., Miller, J.L., Young, N.S., and Liu, J.M. Phenotypic correction of fanconi anemia in human hematopoietic cells with a recombinant adeno-associated virus vector. *J. Clin. Invest.* 94, 1440-1448 (1994).
- Wang X., Ponnazhagan, S., Srivastava, A. Adeno-associated virus type 2 DNA replication in vivo: mutation analyses of the D sequence in viral inverted terminal repeats. *J. Virol.* 71, 3077-3082 (1997).
- West, S.C. Processing of recombination intermediates by the RuvABC proteins. *Annu. Rev. Genet.* 31:213-244 (1997).
- Westfall, T.D., Kennedy, C. & Sneddon, P. The ecto-ATPase inhibitor ARL 67156 enhances parasympathetic neurotransmission in the guinea-pig urinary bladder. *Eur. J. Pharmacol.* 329, 169-173 (1997).
- Wu, P., Phillips, M.I., Bui, J. & Terwilliger, E.F. Adeno-associated virus vector-mediated transgene integration into neurons and other nondividing cell targets. *J. Virol.* 72, 5919-5926 (1998).
- Xiao, X., Li, J. & Samulski, R.J. Efficient long-term gene transfer into muscle tissue of immunocompetent mice by adeno-associated virus vector. *J. Virol.* 70, 8098-8108 (1996).
- Yang, C.C. et al. Cellular recombination pathways and viral terminal repeat hairpin structures are sufficient for adeno-associated virus integration in

vivo and *in vitro*. *J. Virol.* 71, 9231-9247 (1997).

Zhang, Y., Jiang, Q., Dudus, L., Yankaskas, J.R., & Engelhardt, J.F. Vector-specific complementation profiles of two independent primary defects in cystic fibrosis airways. *Hum. Gene Ther.* 9, 635-48 (1998).

5

All publications, patents and patent applications are incorporated herein by reference. While in the foregoing specification this invention has been described in relation to certain preferred embodiments thereof, and many details have been set forth for purposes of illustration, it will be apparent to those skilled
10 in the art that the invention is susceptible to additional embodiments and that certain of the details described herein may be varied considerably without departing from the basic principles of the invention.

WHAT IS CLAIMED IS:

1. A composition comprising at least two recombinant adeno-associated viruses (AAV), comprising:
 - a) a first recombinant AAV comprising a first recombinant DNA molecule comprising linked:
 - i) a first DNA segment comprising a 5'-inverted terminal repeat of AAV;
 - ii) a second DNA segment which does not comprise AAV sequences; and
 - iii) a third DNA segment comprising a 3'-inverted terminal repeat of AAV; and
 - b) a second recombinant AAV comprising a second recombinant DNA molecule comprising linked:
 - i) a first DNA segment comprising a 5'-inverted terminal repeat of AAV;
 - ii) a second DNA segment which does not comprise AAV sequences and which second DNA segment is different than the second DNA segment of the first recombinant DNA molecule; and
 - iii) a third DNA segment comprising a 3'-inverted terminal repeat of AAV.
2. The composition of claim 1 further comprising a delivery vehicle.
3. The composition of claim 2 where the vehicle is a pharmaceutically acceptable carrier.
4. The composition of claim 1 wherein the second DNA segment of the first recombinant DNA molecule comprises a portion of an open reading frame operably linked to a promoter.

5. The composition of claim 4 wherein the first recombinant DNA molecule comprises a splice donor site 3' to the open reading frame.
6. The composition of claim 4 wherein the second DNA segment of the second recombinant DNA molecule comprises the remainder of the open reading frame which together with the second DNA segment of the first recombinant DNA molecule encodes a full-length polypeptide.
7. The composition of claim 6 wherein the second DNA segment of the second recombinant DNA molecule comprises a splice acceptor site 5' to the remainder of the open reading frame.
8. The composition of claim 1 wherein the second DNA segment of the first recombinant DNA molecule comprises an enhancer.
9. The composition of claim 1 wherein the second DNA segment of the first recombinant DNA molecule comprises a promoter.
10. The composition of claim 8 or 9 wherein the second DNA segment of the second recombinant DNA molecule comprises at least a portion of an open reading frame.
11. The composition of claim 10 wherein the second DNA segment of the second recombinant DNA molecule further comprises a promoter operably linked to the open reading frame.
12. The composition of claim 1 wherein the second DNA segment of the first recombinant DNA molecule comprises an origin of replication functional in a host cell.
13. The composition of claim 12 wherein the origin is a viral origin of replication.

14. The composition of claim 13 wherein the origin is functional in a human cell.
15. The composition of claim 12 wherein the second DNA segment of the first recombinant DNA molecule further comprises DNA encoding a protein that binds to the origin of replication.
16. The composition of claim 15 wherein the second DNA segment in the second recombinant DNA molecule comprises at least a portion of an open reading frame.
17. The composition of claim 16 wherein the second DNA segment in the second recombinant DNA molecule further comprises a promoter operably linked to the open reading frame.
18. A recombinant adeno-associated viral vector comprising a heterologous origin of replication.
19. A recombinant adeno-associated viral vector comprising at least one heterologous transcriptional regulatory element functional in a host cell and which does not comprise sequences which encode a protein.
20. The vector of claim 19 wherein the element is a promoter.
21. The vector of claim 19 wherein the element is an enhancer.
22. A recombinant adeno-associated viral vector comprising a DNA sequence encoding a protein that binds to a heterologous origin of replication.
23. A plasmid comprising the vector of claim 18, 19 or 22.
24. A host cell contacted with the composition of claim 1.

25. A host cell contacted with at least two recombinant AAV, wherein a first recombinant AAV comprises a first recombinant DNA molecule comprising linked:
- i) a first DNA segment comprising a 5'-inverted terminal repeat of AAV;
 - ii) a second DNA segment which does not comprise adeno-associated viral sequences; and
 - iii) a third DNA segment comprising a 3'-inverted terminal repeat of AAV; and
- wherein a second recombinant AAV comprises a second recombinant DNA molecule comprising linked:
- i) a first DNA segment comprising a 5'-inverted terminal repeat of AAV;
 - ii) a second DNA segment which does not comprise adeno-associated viral sequences and which second DNA segment is different than the second DNA segment of the first recombinant DNA molecule; and
 - iii) a third DNA segment comprising a 3'-inverted terminal repeat of AAV.

26. A method to transfer recombinant DNAs to a host cell, comprising: contacting the host cell with at least two recombinant AAV, wherein a first recombinant AAV comprises a first recombinant DNA molecule comprising linked:
- i) a first DNA segment comprising a 5'-inverted terminal repeat of AAV;
 - ii) a second DNA segment which does not comprise adeno-associated viral sequences; and
 - iii) a third DNA segment comprising a 3'-inverted terminal repeat of AAV; and
- wherein a second recombinant AAV comprises a second recombinant DNA molecule comprising linked:
- i) a first DNA segment comprising a 5'-inverted terminal

- repeat of AAV;
 - ii) a second DNA segment which does not comprise adeno-associated viral sequences and which second DNA segment is different than the second DNA segment of the first recombinant DNA molecule; and
 - iii) a third DNA segment comprising a 3'-inverted terminal repeat of AAV
27. A method to transfer and express a polypeptide in a host cell comprising contacting the host cell with the composition of claim 1.
28. The method of claim 26 or 27 wherein the second DNA segment of the first recombinant DNA molecule comprises a portion of an open reading frame operably linked to a promoter.
29. The method of claim 28 wherein the first recombinant DNA molecule comprises a splice donor site 3' to the open reading frame.
30. The method of claim 29 wherein the second DNA segment of the second recombinant DNA molecule comprises the remainder of the open reading frame which together with the second DNA segment of the first recombinant DNA molecule encodes a full-length polypeptide.
31. The method of claim 30 wherein the second DNA segment of the second recombinant DNA molecule comprises a splice acceptor site 5' to the remainder of the open reading frame.
32. The method of claim 26 or 27 wherein the second DNA segment of the first recombinant DNA molecule comprises an enhancer.
33. The method of claim 26 or 27 wherein the second DNA segment of the first recombinant DNA molecule comprises a promoter.

34. The method of claim 32 wherein the second DNA segment of the second recombinant DNA molecule comprises at least a portion of an open reading frame.
35. The method of claim 33 wherein the second DNA segment of the second recombinant DNA molecule comprises at least a portion of an open reading frame.
36. The method of claim 34 wherein the second DNA segment of the second recombinant DNA molecule further comprises a promoter operably linked to the open reading frame.
37. The method of claim 35 wherein the second DNA segment of the second recombinant DNA molecule further comprises a promoter operably linked to the open reading frame.
38. The method of claim 26 or 27 wherein the second DNA segment of the first recombinant DNA molecule comprises an origin of replication functional in a host cell.
39. The method of claim 38 wherein the origin is a viral origin of replication.
40. The method of claim 39 wherein the origin is functional in a human cell.
41. The method of claim 26 or 27 wherein the second DNA segment of the first recombinant DNA molecule further comprises DNA encoding a protein that binds to the origin of replication.
42. The method of claim 41 wherein the second DNA segment in the second recombinant DNA molecule comprises a portion of an open reading frame.

43. The method of claim 41 wherein the second DNA segment in the second recombinant DNA molecule further comprises a promoter operably linked to the open reading frame.
44. A recombinant adeno-associated viral vector comprising a DNA sequence encoding a protein that binds to a heterologous origin of replication for use in medical therapy.
45. A recombinant adeno-associated viral vector comprising a heterologous origin of replication for use in medical therapy.




Figure 1




Figure 2






Figure 3





B



Figure 5




Figure 6




Figure. 7




Figure 8




Figure 9

A

10	20	30	40	50	60
GCATGCAAGC	TGTAGATAAG	TAGCATGGCG	GGTTAATCAT	TAACTACAAG	GAACCCCTAG
CGTACGTTTC	ACATCTATTTC	ATCGTACCGC	CCAATTAGTA	ATTGATGTTC	CTTGGGGGATC
70	80	90	100	110	120
TGATGGAGTT	GGCCACTCCC	TCTCTGCGCG	CTCGCTCGCT	CACTGAGGCC	GGGCGGCCAA
ACTACCTCAA	CCGGTGAGGG	AGAGACGCGC	GAGCGAGCGA	GTGACTCCGG	CCCGCCGGTT
130	140	150	160	170	180
AGGTCGCCCC	ACGCCCCGGC	TTTGCCCCGG	CGGCCTCAGT	GAGCGAGCGA	GCGCGCAGAG
TCCAGCGGGC	TGCGGGCCCC	AAACGGGCC	GCCGGAGTCA	CTCGCTCGCT	CGCGCGTCTC
190	200	210	220	230	240
AGGGAGTGGC	CAACTCCATC	ACTAGGGGTT	CCTTGTAAGT	AATGATTAA	CCGCCATGCT
TCCCTCACCG	GTTGAGGTAG	TGATCCCCAA	GGAACATCAA	TTACTAATTG	GGCGGTACGA
250	260	270	280		
ACTTATCTAC	CGATGAATTC	GAGCTTGCAT	GC.....		
TGAATAGATG	GCTACTTAAG	CTCGAACGTA	CG.....		

B

10	20	30	40	50	60
GCATGCAAGC	TGTAGATAAG	TAGCATGGCG	GGTTAATCAT	TAACTACAAG	GAACCCCTAG
CGTACGTTTC	ACATCTATTTC	ATCGTACCGC	CCAATTAGTA	ATTGATGTTC	CTTGGGGGATC
70	80	90	100	110	120
TGATGGAGTT	GGCCACTCCC	TCTCTGCGCG	CTCGCTCGCT	CACTGAGGCC	GGGCGCGCGC
ACTACCTCAA	CCGGTGAGGG	AGAGACGCGC	GAGCGAGCGA	GTGACTCCGG	CCCGCGCGCG
130	140	150	160	170	180
TGCTCGCTC	ACTGAGGCCG	GGCGACCAAA	GGTCGCCCCA	GCCCCGGGCTT	TGCCCCGGGGC
AGCGAGCGAG	TGACTCCGGC	CCGCTGGTTT	CCAGCGGGCT	CGGGCCCCGAA	ACGGGCCCCG
190	200	210	220	230	240
GCCTCAGTGA	GCGAGCGCGC	GCGCAGAGAG	GGAGTGGCCA	ACTCCATCAC	TAGGGGTTCC
CGGAGTCACT	CGCTCGCGCG	CGCGTCTCTC	CCTCACCAGT	TGAGGTAGTG	ATCCCCAAGG
250	260	270	280	290	300
TTGTAGTTAA	TGATTAAACC	GCCATGCTAC	TTATCTACCG	ATGAATTGCA	GCTTGCATGC
AACATCAATT	ACTAATTGGG	CGGTACGATG	AATAGATGGC	TACTTAAGCT	CGAACGTACG

Figure 10

C

10	20	30	40	50	60
GCATGCAAGC	TGTAGATAAG	TAGCATGGCG	GGTTAATCAT	TAACTACAAG	GAACCCCTAG
CGTACGTTTC	ACATCTATTTC	ATCGTACCGC	CCAATTAGTA	ATTGATGTTC	CTTGGGGATC
70	80	90	100	110	120
TGATGGAGTT	GGCCACTCCC	TCTCTGCGCG	CTCGCTCGCT	CACTGAGGCC	GGGCGACCAA
ACTACCTCAA	CCGGTGAGGG	AGAGACGCGC	GAGCGAGCGA	GTGACTCCGG	CCCGCTGGTT
130	140	150	160	170	180
AGGTGCGCCG	ACGCCCAGGC	TTTGGTGCGC	CGGCCTCAGT	GAGCGAGCGA	GCGCGCAGAG
TCCAGCGGGC	TGCGGGCCCC	AAACCAGCGG	GCCGGAGTCA	CTCGCTCGCT	CGCGCGTCTC
190	200	210	220	230	240
AGGGAGTGGC	CAACTCCATC	ACTAGGGGTT	CCTTGTTAGTT	AATGATTAAAC	CCGCCATGCT
TCCCTCACC	GTTGAGGTAG	TGATCCCCAA	GGAACATCAA	TTACTAATTG	GGCGGTACGA
250	260	270	280		
ACTTATCTAC	CGATGAATTC	GAGCTTGCAAT	GC.....		
TGAATAGATG	GCTACTTAAG	CTCGAACGTA	CG.....		

		10	20	30	40	50	
P81	1	GCATGCAAGC	TGTAGATAAG	TAGCATGGCG	GGTTAATCAT	TAACTACAAG	50
p79	1	GCATGCAAGC	TGTAGATAAG	TAGCATGGCG	GGTTAATCAT	TAACTACAAG	50
p1202	1	GCATGCAAGC	TGTAGATAAG	TAGCATGGCG	GGTTAATCAT	TAACTACAAG	50
		60	70	80	90	100	
P81	51	GAACCCCTAG	TGATGGAGTT	GGCCACTCCC	TCTCTGCGCG	CTCGCTCGCT	100
p79	51	GAACCCCTAG	TGATGGAGTT	GGCCACTCCC	TCTCTGCGCG	CTCGCTCGCT	100
p1202	51	GAACCCCTAG	TGATGGAGTT	GGCCACTCCC	TCTCTGCGCG	CTCGCTCGCT	100
		110	120	130	140	150	
P81	101	CACTGAGGCC	GGGCG-----	-----	-----	---GCCAAAG	150
p79	101	CACTGAGGCC	GGGCGCGCGC	TCGCTCGCTC	ACTGAGGCCG	GGCGaCcaAa	150
p1202	101	CACTGAGGCC	GGGCG-----	-----	-----	---ACCAAAG	150
		160	170	180	190	200	
P81	151	GTCGCCCCGAC	GCCCCGGGCTT	TGCCCCGGGCG	GCCTCAGTGA	GCGAGCGAGC	200
p79	151	GgtcgCCcga	GCCCCGGGCTT	TGCCCCGGGCG	GCCTCAGTGA	GCGAGCGcGC	200
p1202	151	GTCGCCCCGAC	GCCCCGGGCTT	TGgtCGccCG	GCCTCAGTGA	GCGAGCGAGC	200
		210	220	230	240	250	
P81	201	GCGCAGAGAG	GGAGTGGCCA	ACTCCATCAC	TAGGGGTTCC	TTGTAGTTAA	250
p79	201	GCGCAGAGAG	GGAGTGGCCA	ACTCCATCAC	TAGGGGTTCC	TTGTAGTTAA	250
p1202	201	GCGCAGAGAG	GGAGTGGCCA	ACTCCATCAC	TAGGGGTTCC	TTGTAGTTAA	250
		260	270	280	290	300	
P81	251	TGATTAACCC	GCCATGCTAC	TTATCTACCG	ATGAATTCGA	GCTTGcATGC	300
p79	251	TGATTAACCC	GCCATGCTAC	TTATCTACCG	ATGAATTCGA	GCTTGcATGC	300
p1202	251	TGATTAACCC	GCCATGCTAC	TTATCTACCG	ATGAATTCGA	GCTTGcATGC	300

Figure 11




Figure 12A




Figure 12B




Figure 12C





Figure 13



B




Figure 14




Figure 16







Figure 7



Feenup (8A-0B)







Figure 18 C + D



Flanell (9)




Fig 19C and b




FIG. 20




FIG. 21




FIG. 22





FIG. 23






Fig. 2A

A



B



C




Fig.25




Fig 2b




Fig 27

SEQUENCE LISTING

<110> University of Iowa Research Foundation
5 Engelhardt, John F.
Duan, Dongsheng
Yan, Ziyang

<120> Adeno-associated virus vectors and uses thereof
10

<130> 875.024WO1

<150> US 60/158,209
<151> 1999-10-07
15

<160> 13

<170> FastSEQ for Windows Version 4.0

20<210> 1
<211> 20
<212> DNA
<213> Adeno-associated virus

25<400> 1
cggggggtcgt tgggcggtca 20

<210> 2
<211> 19

30<212> DNA
<213> Adeno-associated virus

<400> 2
gggcggagcc tatggaaaa 19

35

<210> 3
<211> 505
<212> DNA
<213> Artificial Sequence

40<220>
<223> A consensus sequence.

<400> 3

```

cggggggtcgt tgggcggtca gccaggcggg ccatttaccg taagttatgt aacgactgca      60
ggcatgcaag ctcgaattca tcggtagata agtagcatgg cgggttaatc attaactaca      120
aggaaccct agtgatggag ttggccactc cctctctgcg cgctcgctcg ctcgctgagg      180
5ccgggcgacc aaaggtcgcc cgacgcccgg gctttgcccg ggcggcctca gtgagcgagc      240
gagcgcgcag ctgcgcgctc gctcgctcac tgaggccgcc cgggcaaagc cggggcgctg      300
ggcgaccttt ggtcgcccgg ctcagcgag cgagcgagcg cgcagagagg gagtggccaa      360
ctccatcact aggggttcct tgtagttaat gattaaccg ccatgctact tatctacagc      420
ttgcatgcat gtgagcaaaa ggccagcaaa aggccaggaa ccgtaaaaag gccgcgttgc      480
10tggcggtttt ccataggctc cgccc                                     505

```

<210> 4

<211> 272

<212> DNA

15<213> AAV circular intermediate, clone p81

<400> 4

```

gcatgcaagc tgtagataag tagcatggcg ggttaatcat taactacaag gaaccctag      60
tgatggagtt ggccactccc tctctgcgcg ctcgctcgct cactgaggcc gggcgcccaa      120
20aggtcgccc acgcccgggc tttgcccggg cggcctcagt gagcgagcga gcgcgcagag      180
agggagtggc caactccatc actaggggtt ccttgtagtt aatgattaac ccgcatgct      240
acttatctac cgatgaattc gagcttgcag gc                                     272

```

<210> 5

25<211> 300

<212> DNA

<213> AAV circular intermediate, clone p79

<400> 5

```

30gcatgcaagc tgtagataag tagcatggcg ggttaatcat taactacaag gaaccctag      60
tgatggagtt ggccactccc tctctgcgcg ctcgctcgct cactgaggcc gggcgcgcg      120
tcgctcgctc actgaggccg ggcgaccaa ggtcgcccga gcccgggctt tgcccgggcg      180
gcctcagtga gcgagcgcg gcgcagagag ggagtggcca actccatcac taggggttcc      240
ttgtagttaa tgattaacc gccatgctac ttatctaccg atgaattcga gcttgcagtc      300

```

35

<210> 6

<211> 272

<212> DNA

<213> AAV circular intermediate, clone p1202

40

<400> 6

gcatgcaagc ttagataag tagcatggcg ggttaatcat taactacaag gaacccttag 60
tgatggagtt ggccactccc tctctgcgcg ctcgctcgct cactgaggcc gggcgaccaa 120
aggtcgcccc acgcccgggc tttggtcgcc cggcctcagt gagcgagcga gcgcgcagag 180
5agggagtggc caactccatc actaggggtt ccttgtagtt aatgattaac ccgccatgct 240
acttatctac cgatgaattc gagcttgcac gc 272

<210> 7

<211> 165

10<212> DNA

<213> Unknown

<220>

<223> SEQ ID NO:1 of U.S. Patent No. 5,478,745

15

<400> 7

aggaaccctt agtgatggag ttggccactc cctctctgcg cgctcgctcg ctcactgagg 60
ccgggcgacc aaaggtcgcc cgacgcccgg gctttgcccc ggcgggcctca gtgagcgagc 120
gagcgcgag agagggagtg gccaaactcca tcactagggg ttcct 165

20

<210> 8

<211> 282

<212> DNA

<213> rAAV circular intermediate, clone p79

25

<400> 8

ggcgggcat ttaccgtaag ttatgtggcg actgcaggca tgcaagctcg aattcatcgg 60
tagataagta gcatggcggg ttaatcattg cctacaaaga gcccttagtg atggagtggg 120
ccactccctc tcttcgccga gcgcgcagag agggagtggc caactccctc actaggggtt 180
30cctggcagtt aatgattaac ccgccatgct acttatctac agcttgcag catgtgagca 240
aaaggccagc aaaaggccag gaaccgtaaa aaggccgcgt tg 282

<210> 9

<211> 345

35<212> DNA

<213> rAAV circular intermediate, clone p80

<400> 9

ggccatttac cgtaagtatt gtaacgactg caggcatgca agctcgaatt catcggtaga 60
40taagtagcat ggcgggttaa tcattaacta caaggaaccc ctagtgatgg agttggccac 120

```

tccctctctg cgcgctcgct cgctcgctca ggccgggcca ccaaaggctg cccgacgccc      180
gcccggcctc agcgagcgag cgagcgcgca gagagggagt ggccaactcc atcactaggg      240
gttccttgta gttaatgatt aaccggccat gctacttatt tacagcttgc atgcatgtga      300
gcaaaaaggcc agcaaaaaggc caggaaccgt aaaaaggccg cgttg                      345

```

5

<210> 10

<211> 276

<212> DNA

<213> rAAV circular intermediate, clone p81

10

<400> 10

```

ggccatttac cgtaagttat gtggcgactg caggcatgca agctcgaatt catcggtaga      60
taagtagcat ggcggtttaa tcattgccta caaagagccc ctagtgatgg agccggcct      120
caccgagcga gcgagcgcg cagagaggag tggccaactc catcactagg ggttccttgt      180
15agttaatgat taaccgcca tgctacttat ctacagcttg catgcatgtg agcaaaaagg      240
cagcaaaaagg ccaggaaccg taaaaaggcc gcgttg                      276

```

<210> 11

<211> 316

20<212> DNA

<213> rAAV circular intermediate, clone p86

<400> 11

```

ggccatttac cgtaagttat gtaacgactg caggcatgca agctcgaatt catcggtaga      60
25taagtagcat ggcggtttaa tcattaacta caaggaaccc ctagtgatgg agttggccac      120
tccctctctg cgcgctcgct cgctcgctga ggccgccccg gcctcagcga gcgagcgagc      180
gcgcagagag ggactggcca actccatcac taggggttcc ttgtagttaa tgattaaccc      240
gccatgctac ttatctacag cttgcatgca tgtgagcaaa aggccagcaa aaggccagga      300
accgtaaaaa ggccgc                      316

```

30

<210> 12

<211> 208

<212> DNA

<213> rAAV circular intermediate, clone p87

35

<400> 12

```

ggccatttac cgtaagttat gtaacgactg caggcatgca agctcgaatt catcggtaga      60
taagtagcat ggcggttac tcattgccta caaagagccc ctagtgatgg aattggaatg      120
attcaccctc catgctactt atctacagct tgcattgatg tgagcaaaaag gccagcaaaa      180
40ggccaggaac cgtaaaaagg ccgcgttg                      208

```

<210> 13

<211> 310

<212> DNA

<213> rAAV circular intermediate, clone p88

5

<400> 13

gccatttacc gtaagttatg taacgactgc aggcattgcaa gctcgaattc atcggtagat	60
aagtagcatg gcgggttaat cattgcctac aaagagcccc tagtgatgga gttggccact	120
ccctctctgc gcgctcgctc gctgggcccc gcctcagcga gcgagcgagc gcgcagagag	180
10ggagtggcca actccatcac taggggttcc ttgtagttaa tgattaacct gccatgctac	240
ttatctacag cttgcatgca tgtgagcaaa aggccagcaa aaggccagga accgtaaaaa	300
ggccgcgttg	310

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/864 C12N 01 C12N5/10 A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X, L	WO 99 60146 A (ENGELHARDT JOHN F ;UNIV IOWA RES FOUND (US); DUAN DONGSHENG (US)) 25 November 1999 (1999-11-25) the whole document L:Priority	1-11, 18-21, 23-40,45
X	DUAN D. ET AL.: "Circular intermediates of recombinant adeno-associated virus have defined structural characteristics responsible for long-term episomal persistence in muscle tissue" JOURNAL OF VIROLOGY, vol. 72, no. 11, November 1998 (1998-11), pages 8568-8577, XP002114764 ISSN: 0022-538X the whole document	18-21,45

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

5 March-2001

Date of mailing of the international search report

19/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Mandl, B

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DUAN D. ET AL.: "Formation of adeno-associated virus circular genomes is differentially regulated by adenovirus E4 ORF6 and E2a gene expression"</p> <p>JOURNAL OF VIROLOGY, vol. 73, no. 1, January 1999 (1999-01), pages 161-169, XP002114765 ISSN: 0022-538X the whole document</p>	18-21,45
X	<p>US 5 691 176 A (MCNALLY MAUREEN A ET AL) 25 November 1997 (1997-11-25) column 2, last paragraph</p>	22,23,44
A	<p>PUTTARAJU M. ET AL.: "Spliceosome-mediated RNA trans-splicing as a tool for gene therapy"</p> <p>NATURE BIOTECHNOLOGY, vol. 17, no. 3, March 1999 (1999-03), pages 246-252, XP002130564 ISSN: 1087-0156 the whole document</p>	1-17, 24-43
A	<p>RAMAGE A. D. ET AL.: "Improved EBV-based shuttle vector system: Dicistronic mRNA couples the synthesis of the Epstein-Barr nuclear antigen-1 protein to neomycin resistance."</p> <p>GENE, vol. 197, no. 1-2, 1997, pages 83-89, XP004126404 ISSN: 0378-1119 the whole document</p>	15,22, 23,44
P,X	<p>YAN Z. ET AL.: "Trans-splicing vectors expand the utility of adeno-associated virus for gene therapy."</p> <p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES, vol. 97, no. 12, 6 June 2000 (2000-06-06), pages 6716-6721, XP002162085 June 6, 2000 ISSN: 0027-8424 the whole document</p>	1-45
P,X	<p>YANG J. ET AL.: "Concatamerization of adeno-associated virus circular genomes occurs through intermolecular recombination."</p> <p>JOURNAL OF VIROLOGY, vol. 73, no. 11, November 1999 (1999-11), pages 9468-9477, XP002162086 ISSN: 0022-538X the whole document</p>	18-21

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9960146	A	25-11-1999	AU	4091299 A	06-12-1999
US 5691176	A	25-11-1997	US	5354678 A	11-10-1994
			US	5173414 A	22-12-1992
			US	5681731 A	28-10-1997
			US	5780280 A	14-07-1998
			US	5589377 A	31-12-1996
			AT	130870 T	15-12-1995
			CA	2054517 A,C	01-05-1992
			DE	69114997 D	11-01-1996
			DE	69114997 T	18-04-1996
			DK	488528 T	27-12-1995
			EP	0488528 A	03-06-1992
			HK	1007765 A	23-04-1999
			JP	2655771 B	24-09-1997
			JP	5308975 A	22-11-1993

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.